



Net Content.....



10% HEPTACHLOR GRANULES

CONTROLS:

| | |
|--------------------|---------|
| ACTIVE INGREDIENTS | |
| Heptachlor* | 10.00% |
| Related Compounds | 3.88% |
| INERT INGREDIENTS | 86.12% |
| | <hr/> |
| | 100.00% |

*Heptachlorotetrahydro-4, 7-methanoindene

DIRECTIONS FOR USE

This product is limited strictly to use on lawns and ornamentals so as to avoid contamination of food and feed crops.

BROADCAST TREATMENT: For control of **Ants, Japanese Beetle Larvae, White Grubs, White Fringed Beetle Larvae, European Chafer Larvae** and **Asiatic Garden Beetle Larvae**, apply 20 lbs. of 10% Heptachlor Granules per acre or 1 2 lb. per 1000 sq. ft. to non-food crops including lawns, cemeteries, golf courses, park areas and ornamental nurseries. Wash into top few inches of soil.

For control of **Chiggers, Fleas, Mosquitos and Ticks**, apply 10 lbs. of 10% Heptachlor Granules per acre or 1 1 lb. per 1000 sq. ft. to the above-mentioned areas.

MOUND TREATMENT OF IMPORTED FIRE ANT: Apply 1 or 2 cups of 10% Heptachlor Granules to the entrance of the nest and an area 3 feet around it. If necessary, repeat treatment 2 weeks after first application. Treatment must be limited to non-food crop areas.

WARNING

KEEP OUT OF REACH OF CHILDREN

MAY BE FATAL IF SWALLOWED. DO NOT BREATHE VAPOR, DUST OR SPRAY MIST. DO NOT GET ON SKIN. IN CASE OF CONTACT, WASH IMMEDIATELY WITH SOAP AND WATER. DO NOT CONTAMINATE FOOD AND FEEDSTUFFS.

ANTIDOTE

EXTERNAL—In case of spillage on skin, wash promptly with soap and water.

INTERNAL—Emetic of mustard (1 tablespoon) in tumbler of water. Call a physician.

HIGHLY TOXIC TO FISH AND WILDLIFE. DO NOT CONTAMINATE ANY BODY OF WATER BY DIRECT APPLICATION, CLEANING OF EQUIPMENT OR DISPOSAL OF WASTES AND CONTAINERS. NOTE: BIRDS FEEDING OVER AREAS MAY BE KILLED.

1M 657

STEPHENSON CHEMICAL CO. INC.
COLLEGE PARK, GEORGIA