

DIRECTIONS

FOR USE BY PROFESSIONAL

(A) For control of Norway rats, roof rats, and mice in and around buildings, use Zinc Phosphide at 1% by weight in a suitable grain mixture. Mix at a rate of 1 ounce Zinc Phosphide with six (6) pounds of grain.

Suspend the Zinc Phosphide in four times its weight or volume of corn oil, cottonseed oil, light mineral oil or warm bacon fat, and add to the grain, mixing thoroughly. Store in air-tight containers.

Place bait in tamper-proof boxes made of wood or metal with baffled entrances. Place bait boxes along walls, in corners, and in concealed locations where rats and mice can feed unobserved.



Collect and bury or burn uneaten baits after 24 hours. Do not re-treat the same area with Zinc Phosphide in less than 30-day intervals.

(B) For control of rats around dumps use a 2% Zinc Phosphide bait on grain or meat. Mix at a rate of 2 ounces Zinc Phosphide with six (6) pounds of bait material.

For grain baits mix as directed in part (A) above. For meat baits blend 1 part Zinc Phosphide with 4 parts flour by volume. Add glycerine oil or water to make a paste. Use as a spread on raw meat or mix thoroughly with hamburger or fish. In hot weather add 1/10 of 1% Sodium Sulfite as preservative. When making baits of this type re-

(CONTINUED ON OTHER SIDE PANEL)

EPA Reg. No. 4887-68



ZINC PHOSP

DANGER

KEEP OUT OF
(SEE ANTIDOTE AN
AN EFFECT

CONTROLS: OF NORWA
HOUSE MI

ACTIVE INGREDIENT:
*Zinc Phosphide _____
INERT INGREDIENTS:
Not more than _____
*Expressed as metallic zin

Net Weight
OPERATORS ONLY



ST

OSPHIDE (RUMETAN)

DANGER  **POISON**

OUT OF REACH OF CHILDREN
(SEE AND WARNING ON SIDE PANEL)
EFFECTIVE POISON FOR CONTROL
OF BURROW RATS, ROOF RATS AND
HOUSE MICE

DIRECTIONS

(CONTINUED FROM OTHER SIDE PANEL)

Remember that dogs and cats will actively search out fish and meat baits. Care must be taken to prevent feeding by these animals. Place teaspoon quantities of bait at 8 to 10 foot intervals where rats feed, water, or travel.

Collect and burn or bury uneaten baits after 24 hours. Do not re-treat the same area with Zinc Phosphide in less than 30-day intervals.

DO NOT USE FOODS FOR BAIT WHICH WILL BE ATTRACTIVE TO CHILDREN, SUCH AS COOKIES, CANDY, NUTS, ETC.

DANGER

Baits should be placed in areas not accessible to children, pets, wildlife and domestic animals or in tamper-proof bait boxes.

This product is toxic to birds and other wildlife. Treated bait exposed on soil surface may be hazardous to birds and other wildlife. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

CONTAINER DISPOSAL: Do not reuse empty container. Destroy it by perforating and crushing. Bury in a safe place away from water supplies.

ANTIDOTE

ANTIDOTE: Give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Have victim lie down and keep warm. Call a Physician immediately!

INGREDIENT:		
.....	Not less than	94%
INGREDIENTS:		
.....		6%
metallic zinc - 76%	Total	100%

ATED

3-1975

INSECTICID
ENTICIDE AC
ON REGISTER
MENTS:



ZINC PHOSPHIDE (RUMETAN)

DANGER  POISON

**KEEP OUT OF REACH OF CHILDREN
(SEE ANTIDOTE AND WARNING ON SIDE PANEL)
AN EFFECTIVE POISON FOR CONTROL**

**OLS: OF NORWAY RATS, ROOF RATS AND
HOUSE MICE**

ACTIVE INGREDIENT:		
*Zinc Phosphide	Not less than	94%
INERT INGREDIENTS:		
Not more than		6%
*Expressed as metallic zinc - 76%	Total	100%

DIRECTIONS

(A) For control of Norway rats, roof rats, and mice in an area use a mixture of Zinc Phosphide at 1% by weight in a suitable grain mixture. Mix 1 part Zinc Phosphide with six (6) pounds of grain.

Suspend the Zinc Phosphide in four times its weight or volume of kerosene oil, light mineral oil or warm bacon fat, and add to the grain mixture. Store in air-tight containers.

Place bait in tamper-proof boxes made of wood or metal. Distribute bait boxes along walls, in corners, and in concealed locations where rats feed unobserved.

Collect and bury or burn uneaten baits after 24 hours. Distribute baits with Zinc Phosphide in less than 30-day intervals.

(B) For control of rats around dumps use a 2% Zinc Phosphide mixture. Mix at a rate of 2 ounces Zinc Phosphide with six (6) pounds of grain. For grain baits mix as directed in part (A) above. For meat baits mix Zinc Phosphide with 4 parts flour by volume. Add glycerine or kerosene oil. Use as a spread on raw meat or mix thoroughly with ham or sausage. Always add 1/10 of 1% Sodium Sulfite as preservative. When used on meat.

(CONTINUED ON OTHER SIDE PAGE)

EPA Reg. No. 4887-68



Net Weight _____

IONAL PEST CONTROL OPERATORS ONLY



ZINC PHOSPHIDE (RUMETAN)

DANGER  **POISON**

KEEP OUT OF REACH OF CHILDREN
(SEE INSTRUCTIONS AND WARNING ON SIDE PANEL)
EFFECTIVE POISON FOR CONTROL
NORWAY RATS, ROOF RATS AND
HOUSE MICE

INGREDIENTS:		
Phosphide	Not less than	94%
ADJUVANTS:		
Rumetan		6%
metallic zinc - 76%	Total	100%

Manufactured by
STEPHENSON CHEMICAL CO., INC.
COLLEGE PARK, GEORGIA

DIRECTIONS

(CONTINUED FROM OTHER SIDE PANEL)

Remember that dogs and cats will actively search out fish and meat baits. Care must be taken to prevent feeding by these animals. Place teaspoon quantities of bait at 8 to 10 foot intervals where rats feed, water, or travel.

Collect and burn or bury uneaten baits after 24 hours. Do not re-treat the same area with Zinc Phosphide in less than 30-day intervals.

DO NOT USE FOODS FOR BAITS WHICH WILL BE ATTRACTIVE TO CHILDREN, SUCH AS COOKIES, CANDY, NUTS, ETC.

DANGER

Baits should be placed in areas not accessible to children, pets, wildlife and domestic animals or in tamper-proof bait boxes.

This product is toxic to birds and other wildlife. Treated bait exposed on soil surface may be hazardous to birds and other wildlife. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

CONTAINER DISPOSAL: Do not reuse empty container. Destroy it by perforating or crushing. Bury in a safe place away from water supplies.

ANTIDOTE

ANTIDOTE: Give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Have victim lie down and keep warm. Call a Physician immediately!

DIRECTIONS

(A) For control of Norway rats, roof rats, and mice in and around buildings, use Zinc Phosphide at 1% by weight in a suitable grain mixture. Mix at a rate of 1 ounce Zinc Phosphide with six (6) pounds of grain.

Suspend the Zinc Phosphide in four times its weight or volume of corn oil, cottonseed oil, light mineral oil or warm bacon fat, and add to the grain, mixing thoroughly. Store in air-tight containers.

Place bait in tamper-proof boxes made of wood or metal with baffled entrances. Place bait boxes along walls, in corners, and in concealed locations where rats and mice can feed unobserved.

Collect and bury or burn uneaten baits after 24 hours. Do not re-treat the same area with Zinc Phosphide in less than 30-day intervals.

(B) For control of rats around dumps use a 2% Zinc Phosphide bait on grain or meat. Mix at a rate of 2 ounces Zinc Phosphide with six (6) pounds of bait material.

For grain baits mix as directed in part (A) above. For meat baits blend 1 part Zinc Phosphide with 4 parts flour by volume. Add glycerine oil or water to make a paste. Use as a spread on raw meat or mix thoroughly with hamburger or fish. In hot weather add 1/10 of 1% Sodium Sulfite as preservative. When making baits of this type re-

(CONTINUED ON OTHER SIDE PANEL)

EPA Reg. No. 4887-68

Net Weight _____
FOR USE BY PROFESSIONAL PEST CONTROL OPERATORS OF

ACCEPTED

July 23-1975

4887-68



ZINC PHOSPHIDE (RUMETAN)

DANGER  POISON

KEEP OUT OF REACH OF CHILDREN
(SEE ANTIDOTE AND WARNING ON SIDE PANEL)
AN EFFECTIVE POISON FOR CONTROL

CONTROLS: OF NORWAY RATS, ROOF RATS AND HOUSE MICE

ACTIVE INGREDIENT:		
*Zinc Phosphide	Not less than	94%
INERT INGREDIENTS:		
Not more than		6%
*Expressed as metallic zinc - 76%	Total	100%

Manufactured by
STEPHENSON CHEMICAL CO., INC.
COLLEGE PARK, GEORGIA

Net Weight _____

BY PROFESSIONAL PEST CONTROL OPERATORS ONLY

TED

3-1979

87-68



ZINC PHOSPHIDE (RUMETAN)

DANGER  POISON

KEEP OUT OF REACH OF CHILDREN
(SEE ANTIDOTE AND WARNING ON SIDE PANEL)

**OLS: AN EFFECTIVE POISON FOR CONTROL
OF NORWAY RATS, ROOF RATS AND
HOUSE MICE**

DIRECTIONS

(CONTINUED FROM OTHER SIDE PANEL)

member that dogs and cats will actively search out fish and meat baits. Care must be taken to prevent feeding by these animals. Place teaspoon quantities of bait at 8 to 10 foot intervals where rats feed, water, or travel. Collect and burn or bury uneaten baits after 24 hours. Do not re-treat the same area with Zinc Phosphide in less than 30-day intervals.

DO NOT USE FOODS FOR BAITS WHICH WILL BE ATTRACTIVE TO CHILDREN, SUCH AS COOKIES, CANDY, NUTS, ETC.

DANGER

Baits should be placed in areas not accessible to children, pets, wildlife and domestic animals or in tamper-proof bait boxes.

This product is toxic to birds and other wildlife. Treated bait exposed on soil surface may be hazardous to birds and other wildlife. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

CONTAINER DISPOSAL: Do not reuse empty container. Destroy it by perforating or crushing. Bury in a safe place away from water supplies.

ANTIDOTE

ANTIDOTE: Give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Have victim lie down and keep warm. Call a Physician immediately!

ACTIVE INGREDIENT:		
*Zinc Phosphide	Not less than	94%
INERT INGREDIENTS:		
Not more than		6%
*Expressed as metallic zinc - 76% Total		100%

Manufactured by
STEPHENSON CHEMICAL CO., INC.
COLLEGE PARK, GEORGIA