

**DIRECTIONS**

Apply when air is calm, using ground equipment, aircraft, or hand duster. Unless otherwise specified, apply at the first sign of infestation and repeat at 7-10 day intervals when needed to maintain control. Observe use limits given for specific crops. See NOTE below on very small-scale applications.

**BEANS:** Leafhoppers — Use 20 lb. per acre. Lygus Bug, Stink Bugs — Use 40 lb. per acre. Mexican Bean Beetle — Use 10 lb. per acre. Apply thoroughly to upper and lower leaf surfaces. Do not feed treated vines to dairy animals or animals being finished for slaughter.

**CORN:** Corn Earworm — Apply 30-40 lb. per acre starting when the first silks appear. Repeat at 2-3 day intervals throughout the silking period. Direct duster outlets toward ear zone to insure thorough coverage of each silk. May be applied up to harvest on sweet corn ears for human consumption. To avoid excessive residues on fodder and forage, do not apply within 7 days of harvest.

**COTTON:** Early-season — Cotton Fleahopper, Cotton Leafworm, Flea Beetles, Thrips. Use 10 to 20 lb. per acre. Mid- and Late-Season Control — Boll Weevil, Bollworm, Fall Armyworm, Leafhoppers, Leaf Rollers, Tarnished Plant Bug. Use 20 to 40 lb. per acre, depending on size of cotton and insect population level. Applications for Boll Weevil should be made every 5-7 days as long as control is necessary. Pink Bollworm — Use 30 to 60 lb. per acre on a 5-7 day schedule, depending on size of cotton and degree of infestation. Stink Bugs — Use 40 lb. per acre. Do not graze treated fields or feed trash to dairy or meat animals.

**CUCUMBER, SUMMER SQUASH:** Cucumber Beetles, Flea Beetles, Melonworm, Pickleworm, Squash Bugs — Use 20 lbs. per acre.

**EGGPLANT, PEPPER, TOMATO:** Fall Armyworm Flea Beetles, Tomato Fruitworm — Use 20-40 lb. per acre. European Corn Borer, Stink Bugs, Tomato Hornworm — Use 40-80 lb. per acre.

**POTATO:** Colorado Potato Beetle — Use 10 lb. per acre. Flea Beetles, Leafhoppers — Use 20 lb. per acre. Lygus Bugs — Use 40 lb. per acre. Apply dust thoroughly to upper and lower surfaces of leaves. Do not graze treated fields or feed trash to dairy or meat animals.

**NOTE:** On very small-scale applications, disregard the above rates and apply the dust to uniformly cover upper and lower leaf surfaces.



**OBSERVE THE FOLLOWING USE PRECAUTIONS**

Sevin insecticide will not control insects if used with lime or other highly alkaline materials. This product is highly toxic to bees exposed to direct treatment or residues on crops. Do not apply during bloom when honey bees or other pollinators are active. Protective information may be obtained from your Cooperative Agricultural Extension Service. Avoid drift to adjoining food and forage crops.

The Federal Food and Drug Administration has set official tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates, and do not apply closer to harvest than specified.

Consult State Agricultural Experiment Station or State Extension Service Specialists for additional information as the timing, number, and rate of applications needed will vary with local conditions.

Net Wt. 1 Lb.



**5% SEVIN\***  
**DUST**

**CONTROLS:**

<b>ACTIVE INGREDIENTS</b>		
Carbaryl (1-naphthyl N-methylcarbamate)	.....	5%
<b>INERT INGREDIENTS</b>	.....	95%
		<hr/>
	TOTAL.....	100%

\*Trade-Mark of Union Carbide Corp. for 1-naphthyl N-methylcarbamate

Manufactured by  
**STEPHENSON CHEMICAL CO., INC.**  
COLLEGE PARK, GEORGIA

Manufactured by  
**STEPHENSON CHEMICAL CO., INC.**  
COLLEGE PARK, GEORGIA

