

Best copy available

4887-83

05/11/1962

Date Letter APR 13 1962

1/2

Net Wt. _____

STEPHENSON
Chemicals

FLY BAIT

The New Sugar Base Bait with
Instant Acting DDVP!

ACTIVE INGREDIENT

Dimethyl Dichlorovinyl Phosphate----- .5%

INERT INGREDIENTS-----99.5%

100.0%

ACCEPTED

MAY 1 1 1962

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 4887-83

KILLS RESISTANT FLIES!

- Sprinkle as a dry bait
- or
- Mix with water to make liquid bait
- Wet or dry-flies love it, and it kills quickly!

USE AROUND DAIRY BARNs, POULTRY HOUSES, HOG FARMS and OTHER
FAIR APPLICATIONS. NOT RECOMMENDED FOR USE IN HOMES

DIRECTIONS FOR USE

DRY BAIT

As a dry bait, simply shake right out of the can. Scatter lightly on floors, window sills and other areas where flies congregate. Use about 1/4 ounces per 1,000 square feet. A faster kill may be achieved if applied to a moistened surface.

WET BAIT

As a liquid bait, dissolve one pound of dry bait in a half-gallon of water. The resulting liquid may then be sprayed on walls, painted on sills, or used to make bait stations. Bait stations need not be elaborate. Simply place 1/4 inch squares of paper toweling, burlap, cotton, or other absorbent material in a paper plate, pie plate or on a board. Then soak with fly bait. Bait stations should be placed at about 15 ft. intervals in each heavily infested building. After flies have been brought into control, treatment should be repeated at weekly intervals, or when needed.

Important: Do not over-use! Whether used as a liquid or a dry bait, this material is very effective, and a lightly scattered streak or a small bait station will usually be effective.

Economy hint: When liquid bait stations become dry, try dampening them with plain water. Usually this will revive the bait and make it usable another day.

CAUTION: HARMFUL IF SWALLOWED, OR ABSORBED THROUGH THE SKIN. DO NOT CONTAMINATE FEED OR FOODSTUFFS. DO NOT APPLY TO ANIMALS. IF BROUGHT INTO CONTACT WITH THE SKIN, WASH IMMEDIATELY WITH SOAP AND WATER. USE AS A COARSE WET SPRAY. AVOID BREATHING SPRAY MIST. KEEP OUT OF REACH OF CHILDREN AND PETS. DO NOT SCATTER WHERE POULTRY FEED OR WHERE ANIMALS CAN LICK DRY BAIT OR SPRAYED SURFACES. NOT FOR USE IN HOMES OR WHERE MILK IS STORED OR PROCESSED. ATROPINE SULFATE IS ANTIDOTAL.

NOTICE: Having no control over the use of this material, Stephenson Chemical Co., Inc. makes no warranty concerning this material except that it conforms to the chemical content as shown on the label. Neither Stephenson Chemical Co., Inc. nor the seller shall be held responsible in any manner for any personal injury or property damage or other type of loss resulting from handling storage or use of this material. The buyer assumes all risk and liability therefrom and accepts and uses this material on these conditions.

Manufactured by
STEPHENSON CHEMICAL CO., INC.
P. O. Box 188
College Park, Georgia

ACCEPTED
5-11-62
4887-83
UNDER THE FEDERAL INSECTICIDE
ACT
REGISTERED
SUBJECT
COMMENTS

