

USDA Reg. No.
CAUTION 4887-83

HARMFUL IF SWALLOWED, OR ABSORBED THROUGH THE SKIN. DO NOT CONTAMINATE FEED OR FOODSTUFFS. DO NOT APPLY TO ANIMALS. IF BROUGHT INTO CONTACT WITH THE SKIN, WASH IMMEDIATELY WITH SOAP AND WATER. USE AS A COARSE WET SPRAY. AVOID BREATHING SPRAY MIST. KEEP OUT OF REACH OF CHILDREN AND PETS. DO NOT SCATTER WHERE POULTRY FEED OR WHERE ANIMALS CAN LICK DRY BAIT OR SPRAYED SURFACES. NOT FOR USE IN HOMES OR WHERE MILK IS STORED OR PROCESSED. ATROPINE SULFATE IS ANTIDOTAL.

NOTICE: Having no control over the use of this material, Stephenson Chemical Co., Inc. makes no warranty concerning this material except that it conforms to the chemical content as shown on the label. Neither Stephenson Chemical Co., Inc. nor the seller shall be held responsible in any manner for any personal injury or property damage or other type of loss resulting from handling, storage or use of this material. The buyer assumes all risk and liability therefrom and accepts and uses this material on these conditions.

FLY BAIT

ACTIVE INGREDIENT

Dimethyl Dichlorovinyl Phosphate	-----	.5%
INERT INGREDIENTS	-----	99.5%
		100.0%

USE AROUND DAIRY BARN'S
POULTRY HOUSES, HOG FARMS
OTHER FARM APPLICATIONS

Manufactured by
STEPHENSON CHEMICAL CO., INC.
P. O. Box 188, College Park, Ga.

CAUTION ← 14 pt. type
KEEP OUT OF REACH OF CHILDREN
SEE CAUTIONS ON SIDE PANELS

DIRECTIONS FOR USE

DRY BAIT - As a dry bait, simply shake right out of the can. Scatter lightly on floors, window sills and other areas where flies congregate. Use about 4 ounces per 1,000 square feet. A faster kill may be achieved if applied to a moistened surface.

WET BAIT - As a liquid bait, dissolve one pound of dry bait in a half-gallon of water. The resulting liquid may then be sprayed on walls, painted on sills, or used to make bait stations. Bait stations need not be elaborate. Simply place 4 inch squares of paper toweling, burlap, cotton, or other absorbent material in a paper plate, pie plate or on a board. Then soak with fly bait. Bait stations should be placed at about 15 ft. intervals in each heavily infested building. After flies have been brought into control, treatment should be repeated at weekly intervals, or when needed.

Important! Do not over-use! Whether used as a liquid or a dry bait, this material is very effective, and a lightly scattered stream or a small bait station will be effective.

Economy hint - When liquid bait stations become dry, try dampening them with plain water. Usually this will revive the bait and make it usable another day.

ACCEPTED
4-23-65
4887-83

4887-83

04/23/1965

1/1