

NET WEIGHT 1.5 POUNDS



REGISTERED
4-25-67
4887-83
NUMBER THE REGISTERED TRADE MARK OF
FOR ECONOMIC POISON RESISTANCE
ED UNDER NO. 4887-83 SUBJECT
TO ATTACHED COMMENTS.

ACTIVE INGREDIENTS:

*Dichlorovinyl Dimethyl Phosphate	0.47%
*Related Compounds	0.03%
INERT INGREDIENTS	99.50%
	100%

*Equivalent to .50% of Shell Vapona Insecticide

CAUTION: KEEP OUT OF REACH OF CHILDREN (SEE CAUTIONS ON BACK PANEL)

PACKED EXCLUSIVELY FOR
***COOK'S PEST CONTROL, INC.**
DECATUR, ALABAMA

Distributing Offices: Decatur, Ala.—Huntsville, Ala.—Florence, Ala.—Cullman, Ala.
Birmingham, Ala.—Columbia, Tenn.—Tullahoma, Tenn.

- Sprinkle as a dry bait or • Mix with water to make liquid bait
- Wet or dry—flies love it, and it kills quickly!

DIRECTIONS FOR USE

DRY BAIT — As a dry bait, simply shake right out of the can. Scatter lightly on floors, window sills and other areas where flies congregate. Use about 4 ounces per 1,000 square feet. A faster kill may be achieved if applied to a moistened surface.

WET BAIT — As a liquid bait, dissolve one pound of dry bait in a half-gallon of water. The resulting liquid may then be sprayed on walls, painted on sills, or used to make bait stations. Bait stations need not be elaborate. Simply place 4 inch squares of paper toweling, burlap, cotton, or other absorbent material in a paper plate, pie plate or on a board. Then soak with fly bait. Bait stations should be placed at about 15 ft. intervals in each heavily infested building. After flies have been brought into control, treatment should be repeated at weekly intervals, or when needed.

IMPORTANT! DO NOT OVER-USE! Whether used as a liquid or a dry bait, this material is very effective, and a lightly scattered stream or a small bait station will be effective.

ECONOMY HINT—When liquid bait stations become dry, try dampening them with plain water. Usually this will revive the bait and make it usable another day.

CAUTION

HARMFUL IF SWALLOWED, OR ABSORBED THROUGH THE SKIN. DO NOT CONTAMINATE FEED OR FOOD-STUFFS. DO NOT APPLY TO ANIMALS. IF BROUGHT INTO CONTACT WITH THE SKIN, WASH IMMEDIATELY WITH SOAP AND WATER. USE AS A COARSE WET SPRAY. AVOID BREATHING SPRAY MIST. KEEP OUT OF REACH OF CHILDREN AND PETS. DO NOT SCATTER WHERE POULTRY FEED OR WHERE ANIMALS CAN LICK DRY BAIT OR SPRAYED SURFACES. NOT FOR USE IN HOMES OR WHERE MILK IS STORED OR PROCESSED. ATROPINE SULFATE IS ANTIDOTAL. IF SWALLOWED INDUCE VOMITING IMMEDIATELY AND GET MEDICAL ATTENTION.

NOTICE

Having no control over the use of this material, Cook's Pest Control makes no warranty concerning this material except that it conforms to the chemical content as shown on the label. Neither Cook's Pest Control nor the seller shall be held responsible in any manner for any personal injury or property damage or other type of loss resulting from handling, storage or use of this material. The buyer assumes all risk and liability therefrom and accepts and uses this material on these conditions.

USDA Reg. No. 4887-83

4887-83

04/25/1967

1/2

KILLS RESISTANT FLIES!

- Sprinkle as a dry bait
or
- Mix with water to make liquid bait
- Wet or dry-flies love it, and it kills quickly!

Important. Do not over-use! Whether used as a liquid or a dry bait, this material is very effective, and a lightly scattered stream or a small bait station will be effective.

Economy hint—When liquid bait stations become dry, try dampening them with plain water. Usually this will revive the bait and make it useable another day.

NET WT. 1 LB.



FLY BAIT

Sugar Base with DDVP

**USE AROUND DAIRY BARNs, POULTRY
HOUSES, HOG FARMS**

DIRECTIONS

DRY BAIT—As a dry bait, simply shake right out of the can. Scatter lightly on floors, window sills and other areas where flies congregate. Use about 4 ounces per 1,000 square feet. A faster kill may be achieved if applied to a moistened surface.

WET BAIT—As a liquid bait, dissolve one pound of dry bait in a half-gallon of water. The resulting liquid may then be sprayed on walls, painted on sills, or used to make bait stations. Bait stations need not be elaborate. Simply place 4 inch squares of paper toweling, burlap, cotton, or other absorbent material in a paper plate, pie plate or on a board. Then soak with fly bait. Bait stations should be placed at about 15 ft. intervals in each heavily infested building. After flies have been brought into control, treatment should be repeated at weekly intervals, or when needed.

ACTIVE INGREDIENTS	
*Related Compounds	0.03%
*2,2-Dichlorovinyl Dimethyl Phosphate	0.47%
INERT INGREDIENTS	99.50%
TOTAL100.00%	
*Equivalent to .5% by wt. of Shell Vapona	

Manufactured by
STEPHENSON CHEMICAL CO., INC.
COLLEGE PARK, GEORGIA

