


DIRECTIONS

For protection of stored peanuts against infestations of red flour beetle, Indian meal moth, confused flour beetle, rice weevil, flat grain beetle, rusty grain beetle, lesser grain borer, granary weevil, and saw-toothed grain beetle, a complete spraying program must be undertaken using both Stephenson Chemicals Premium Grade 57% Malathion E. C. for Stored Peanuts and Stephenson Chemicals Premium Grade 25% Malathion W. P. for Stored Peanuts following the directions below:

RESIDUAL WAREHOUSE SPRAY — BEFORE STORING PEANUTS: Clean warehouse thoroughly of trash and old remains of peanuts 1 to 2 weeks before new peanut crop is stored. Thoroughly spray with sufficient pressure interior of empty warehouse (including cracks and protected places), outside walls to height of 6 to 8 feet, and the ground to a distance of about 6 feet from warehouse by diluting 1 pint of Stephenson Chemicals Premium Grade 57% Malathion E. C. for stored peanuts in sufficient water to make 2-1,2 gallons of spray or 1 gallon with 19 gallons of water. Apply finished spray at the rate of 2 gallons 1000 square feet or to run-off.

(Continued on other side panel)

NET CONTENTS _____



9-25-72
4887-113

PREMIUM GRADE 57% MALATHION

Emulsifiable Concentrate for Stored Peanuts
USE TO AID IN THE CONTROL OF INSECTS WHICH INFEST STORED PEANUTS.

CAUTION: KEEP OUT OF REACH OF CHILDREN (READ CAUTIONS ON SIDE PANEL)

ACTIVE INGREDIENTS:	
Malathion	57.0%
Xylene	32.0%
INERT INGREDIENTS	
	11.0%
Total	100.0%
O, O-dimethyl dithiophosphate of diethyl mercaptosuccinate.	

Manufactured by
STEPHENSON CHEMICAL CO., INC.
COLLEGE PARK, GEORGIA

DIRECTIONS

For control of stored peanuts against pest insects and fungi, water and oil sprays are not recommended. Rice weevil, flat grain beetle, rusty grain beetle, lesser grain borer, saw-toothed grain beetle, and saw-toothed grain moth. Complete spraying program must be undertaken using both Stephenson Chemicals Premium Grade 57% Malathion E. C. for Stored Peanuts and Stephenson Chemicals Premium Grade 2% Malathion W. P. for Stored Peanuts. Follow the directions below.

RESIDUAL WAREHOUSE SPRAY - BEFORE STORING PEANUTS. Clean warehouse thoroughly of trash and old remains of peanuts 2 to 3 weeks before new peanut crop stored. Thoroughly spray with sufficient pressure interior of empty warehouse including cracks and protected places, outside walls to height of 6 to 5 feet, and the ground to distance of about 6 feet from warehouse by diluting 1 pint of Stephenson Chemicals Premium Grade 57% Malathion E. C. for stored peanuts in sufficient water to make 2 1/2 gallons of spray. Mix 1 gallon with 19 gallons of water. Apply finished spray at the rate of 1 gallon per 1000 feet² to "put off".

Continued on other side page

CAUTION

KEEP OUT OF REACH OF CHILDREN. HARMFUL IF SWALLOWED. FLAMMABLE. AVOID BREATHING SPRAY MIST. AVOID CONTACT WITH SKIN. WASH THOROUGHLY AFTER USE. AVOID CONTAMINATION OF FEED AND FOODSTUFFS. KEEP CONTAINER CLOSED AND AWAY FROM HEAT AND OPEN FLAME.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

NOTE TO PHYSICIAN: Malathion upon use may cause Cholinesterase Inhibition. Atropine is antidotal.

STEPHENSON
Chemicals

PREMIUM GRADE

57% MALATHION

Emulsifiable Concentrate for Stored Peanuts

USE TO AID IN THE CONTROL OF INSECTS WHICH INFEST STORED PEANUTS.

CAUTION: KEEP OUT OF REACH OF CHILDREN. READ CAUTIONS ON SIDE PANEL.

Manufactured by
STEPHENSON CHEMICAL CO., INC.
COLLEGE PARK, GEORGIA

STEPHENSON

Chemicals

PREMIUM GRADE

0% MALATHION

able Concentrate for Stored Peanuts

AID IN THE CONTROL OF INSECTS
INFEST STORED PEANUTS

ION: KEEP OUT OF REACH OF CHILDREN
READ CAUTIONS ON SIDE PANEL

DIRECTIONS (CONTINUED)

BULK SPRAY TREATMENT - PEANUTS CONTAINED IN STEPHENSON CHEMICALS PREMIUM GRADE 2% MALATHION W. P. CONCENTRATE - Apply 1 gallon of water per 15 tons of peanuts, stacked peanuts, or other bulk peanuts, by spraying into bins. Preferably use a suitable mechanical sprayer which will provide uniform application to flow of peanuts. Otherwise, adjust nozzle to spray directly into flow of peanuts.

SUPPLEMENTAL SURFACE SPRAY - After washing bins, clean up spilled peanuts before spraying. Apply surface spray with a sprayer equipped with a nozzle of Premium Grade 2% Malathion W. P. for stored peanuts at a rate of 2 gallons per 1000 square feet of surface. Apply to the surface of bins, if filled and leveled, but not later than the first week of storage. For best results, treatment 7 days later, followed by treatments at 14 and 21 days later. Use a sprayer equipped with a fan blower and nozzle capable of depositing a fine surface application of concentrate. If warm temperatures prevail, increase the treatment interval to 10 days.

CONTAINER DISPOSAL: Do not reuse empty container. Destroy it by perforating or crushing. Bury in a safe place away from water supplies.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

EPA Reg. No. 4887-113

Manufactured by
STEPHENSON CHEMICAL CO., INC.
COLLEGE PARK, GEORGIA