


A 1 lb. quantity of this product is sufficient to control 4 or 5 rats or several mice.



ACCEPTED
6-8-71
with comment
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND FUNGICIDE ACT
FOR ECONOMIC CONTROL
REGISTERED UNDER NO. 4887-100

Net Wt.....



**FUMARIN RAT BAIT
MEAL**

GEN
Place bait in locations where Bait placements along walls where rats and mice can feed be placed in locations not accessible to other animals in tamper-proof containers. If infestation is present (nesting should be established and rats are sour, moldy or otherwise), remove promptly.

Place 1 oz to 1 lb of STEP BAIT where rats are known to feed continuously until complete control is achieved in 14 DAYS. NEVER LET A RAT LIVE MORE THAN 48 HOURS.

M
Use 1 or 2 tablespoons of FUMARIN RAT BAIT MEAL for control of mice. Make sure...

ACTIVE INGREDIENT	
3-(alpha-acetonylfurfuryl)-4-hydroxycoumarin	0.025%
INERT INGREDIENTS	99.975%
	<hr/>
	100.00%

Manufactured by
STEPHENSON CHEMICAL CO., INC
COLLEGE PARK, GEORGIA

ED
7/
INSECTICIDE
ANTHRAQUINONE AC
REGISTERED
87-100

with comment

Net Wt.....



FUMARIN RAT BAIT MEAL

GENERAL DIRECTIONS

Place bait in locations where rats and mice feed, water or travel. Bait placements along walls, in corners, and in concealed locations where rats and mice can feed unobserved are desirable. Bait should be placed in locations not accessible to children, pets, and domestic animals in tamper-proof boxes. Where a continuous source of infestation is present (nearby dumps), permanent bait stations should be established and replenished as needed. NOTE, Baits that are sour, moldy or otherwise decomposed should be replaced promptly.

RAT CONTROL

Place 1 oz to 1 lb of STEPHENSON CHEMICALS FUMARIN RAT BAIT where rats are known to run or feed or water. Baiting must be continuous until complete lack of feeding is noticed, USUALLY 10 to 14 DAYS. NEVER LET A BAIT STATION REMAIN EMPTY MORE THAN 48 HOURS.

MOUSE CONTROL

Use 1 or 2 tablespoons of bait at various 8 to 12 ft. intervals for control of mice. Make sure that bait is available for at least 15 days.

ACTIVE INGREDIENT	
1-(alpha-acetyl-furfuryl)-4-hydroxycoumarin	0.025%
INERT INGREDIENTS	99.975%
	<hr/>
	100.00%

Manufactured by
STEPHENSON CHEMICAL CO., INC.
COLLEGE PARK, GEORGIA

A 1 lb. quantity
of this product
is sufficient to
control 4 or 5
rats or several
mice.



REGISTERED TRADE MARK
PAT. 2,800,000
U.S. PAT. OFF. 11-18-57

Net Wt.

STEPHENSON
Chemicals •

**FUMARIN RAT BAIT
MEAL**
KILLS RATS and MICE

Cheese Odor Added

CAUTION: KEEP OUT OF REACH OF CHILDREN.
(SEE CAUTION ON SIDE PANEL)

CAUTION

THE ACTIVE INGREDIENT, FUMARIN, IS AN ANTI-COAGULANT CHEMICAL WHICH MAY CAUSE SERIOUS HEMORRHAGE IN HUMANS, PETS, OR LIVESTOCK. IF EATEN, GIVE A TABLESPOON OF SALT IN A GLASS OF WARM WATER AND REPEAT UNTIL VOMIT FLUID IS CLEAR. CALL A PHYSICIAN IMMEDIATELY.

NOTE FOR PHYSICIANS

In case of ingestion of FUMARIN, blood transfusions combined with intravenous injections and oral doses of Vitamin K are indicated as in the case of hemorrhage caused by overdoses of Dicumarol.

4887-120

Net Wt.

STEPHENSON

Chemicals

FUMARIN RAT BAIT MEAL

KILLS RATS and MICE

Cheese Odor Added

CAUTION: KEEP OUT OF REACH OF CHILDREN.
(SEE CAUTION ON SIDE PANEL)

ACTIVE INGREDIENT:

1. FUMARIN BAIT MEAL (100%)



NT CHEMI-
NS, PETS,
A GLASS
EAR. CALL

lined with
sted as in

GENERAL DIRECTIONS

Place bait in locations where rats and mice feed, water or travel. Bait placements along walls, in corners, and in concealed locations where rats and mice can feed unobserved are desirable. Bait should be placed in locations not accessible to children, pets, and domestic animals in tamper-proof boxes. Where a continuous source of infestation is present (meatby dumps, permanent bait stations) should be established and replenished as needed. NOTE: Baits that are used in any other wise intended use should be replaced promptly.

RAT CONTROL

Use 1/2 to 1/4 cup of STEPHENSON CHEMICALS FUMARIN RAT BAIT MEAL where rats are known to run or feed or water. Baiting must be continuous until complete lack of feeding is noticed, USUALLY 10 to 14 DAYS. NEVER LET A BAIT STATION REMAIN EMPTY MORE THAN 15 HOURS.

MOUSE CONTROL

Use 1/2 to 1/4 tablespoons of bait at various 8 to 12 ft. intervals for control of mice. Make sure that bait is available for at least 15 days.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

CONTAINER DISPOSAL: Do not reuse empty containers. Crush and bury at least 18 inches deep in a safe place away from water supplies.

STEPHENSON CHEMICALS, INC. No. 4887-120