

NET WT. 1 LB.

May be used on Apples, Apricots, Cherries, Peaches, Pears, Plums and Prunes to control the following insects and diseases

INSECTS: Aphids, many species of mites, codling moth, bud moth, leafrollers, leafhopper, curculio, mealy bug, bagworm, cherry fruit fly and peach twig borer.

DISEASES: Scab, black rot (fruit and leaf spots), bitter rot, brown rot, fly speck, Brooks fruit spot and Botryosphaeria.

DIRECTIONS FOR USE

Use 5 level tablespoons per gallon of water or 1 cup per 2½ gallons of water. Mix well with required amount of water and keep mixture agitated while spraying. The above is 1 pound per 10 gallons of water. Use a pressure sprayer. Spray thoroughly both sides of leaves, branches and trunks. The following table will act as a guide in determining gallons of spray per tree:

Approximate Age of Tree (Years)	Height (Feet)	Spread (Feet)	Gallons To Apply
1-5	5-8	3-6	½-1
5-10	8-12	6-9	1-3
10-15	12-18	9-15	4-8
over 15	18	15	8-10

SUGGESTED SPRAY SCHEDULE

For specific local conditions on timing and number of spray applications consult State Agricultural Experiment Station or State Extension Service authorities.



APPLES AND PEARS: Spray when new spring growth appears and repeat at 7-day intervals up to bloom. After petal fall, repeat at 10-day intervals. Do not apply to apples and pears within 7 days of harvest.

PEACHES: Make first application at pink bud stage. Apply from 5 to 7 sprays at 10-day intervals. Peach Tree Borer—½ lb. in 1 gal. water, spraying or painting trunks of trees from crotch to ground about the first of July, August and September. Do not apply to peaches within 7 days of harvest.

PLUMS AND PRUNES: Make one application just before blossoms open. After petal fall, repeat at 10-day intervals. Apply from 5 to 7 sprays. Do not apply to plums and prunes within 7 days of harvest.

CHERRIES, APRICOTS: Apply at full bloom, petal fall, 2 weeks after petal fall and again 2 weeks before harvest. Do not apply to Cherries and Apricots within 7 days of harvest.

NOTE: THIS PRODUCT WILL KILL FISH. KEEP OUT OF ANY BODY OF WATER. THIS PRODUCT IS HIGHLY TOXIC TO BEES EXPOSED TO DIRECT TREATMENT ON RESIDUES OR CROPS. PROTECTIVE INFORMATION MAY BE OBTAINED FROM YOUR COOPERATIVE AGRICULTURAL EXTENSION SERVICE.



FRUIT TREE SPRAY

Contains Malathion and Captan

CONTROLS:

ACTIVE INGREDIENTS:	
*Malathion	7.5%
**Captan	15.0%
INERT INGREDIENTS: 77.5%	
TOTAL	
	100.0%

*O, O-dimethyl dithiophosphate of diethyl mercaptosuccinate.
**N-trichloromethylthio-4-cyclohexene-1, 2-dicarboximide.

Manufactured by
STEPHENSON CHEMICAL CO., INC.
COLLEGE PARK, GEORGIA

Manufactured by
STEPHENSON CHEMICAL CO., INC.
COLLEGE PARK, GEORGIA

PESTS: Aphids, many species of mites, codling moth, bud moth, leafrollers, leafhopper, and many other insects, including fruit fly and peach twig borer.

DISEASES: Scab, black rot, fruit and leaf spots, bitter rot, brown rot, fly speck, Brooks rot spot, and Botryosphaeria.

DIRECTIONS FOR USE

Apply 1/2 to 1 gallon of spray per acre, depending on tree height. Mix well with water. Spray thoroughly, covering all sides of leaves. Do not apply to fruit or to plants within 7 days of harvest.

Approximate Age of Tree (Years)	Height (Feet)	Sprayer (Feet)	Gallons To Apply
1-5	5-8	3-6	1/2-1
5-10	8-12	6-9	1-3
10-15	12-18	9-15	4-8
over 15	18	15	8-10

SUGGESTED SPRAY SCHEDULE

For specific local conditions on timing and number of spray applications, consult state Agricultural Experiment Station or state Extension Service authorities.

APPLES AND PEARS: Spray when new shoots are with leaves and repeat at 7 day intervals up to bloom. After petal fall repeat at 7 day intervals. Do not apply to apples and pears within 7 days of harvest.

PEACHES: Make first application at first bud break. Repeat 7 to 10 sprays at 10 day intervals. Do not apply to fruit or to plants within 7 days of harvest.

PLUMS AND PRUNES: Spray when new shoots are with leaves and repeat at 7 day intervals up to bloom. Do not apply to fruit or to plants within 7 days of harvest.

CHERRIES: Spray when new shoots are with leaves and repeat at 7 day intervals up to bloom. Do not apply to fruit or to plants within 7 days of harvest.

NOTE: THIS PRODUCT WILL KILL FISH. KEEP OUT OF ANY BODY OF WATER. THIS PRODUCT IS HIGHLY TOXIC TO BEES EXPOSED TO DIRECT TREATMENT ON RESIDUES OR CROPS. PROTECTIVE INFORMATION MAY BE OBTAINED FROM YOUR COOPERATIVE AGRICULTURAL EXTENSION SERVICE.

CAUTION

KEEP OUT OF REACH OF CHILDREN. HARMFUL IF SWALLOWED. AVOID BREATHING OF SPRAY MIST OR DUST. AVOID CONTACT WITH SKIN. WASH THOROUGHLY AFTER USING. AVOID CONTAMINATION OF FEED AND FOODSTUFFS. DO NOT SPRAY FRUIT TREES WHILE IN OPEN BLOOM. WASH ALL FRUIT THOROUGHLY BEFORE USING. DO NOT REUSE EMPTY CONTAINER. DESTROY BY BURYING WITH WASTE.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

USDA Reg. No. 4887-131

STEPHENSON
Chemicals

FRUIT TREE
SPRAY

Contains Malathion and Captan

Meets insects and diseases that appear on Apples, Apricots, Cherries, Peaches, Pears, Plums and Prunes

CAUTION: KEEP OUT OF REACH OF CHILDREN (SEE CAUTION ON BACK PANEL)

ACTIVE INGREDIENTS:
Malathion
Captan
INERT INGREDIENTS

Manufactured by
STEPHENSON CHEMICAL CO., INC.
COLLEGE PARK, GEORGIA

Manufactured by
STEPHENSON CHEMICAL CO., INC.
COLLEGE PARK, GEORGIA