

1M 10-71

TO KILL UNDESIRABLE TREES IN NON-CROPPED AREAS: Make a continuous cut or "frill" around the base of the tree with downward axe strokes, cutting into the sapwood.

RATE: Apply dry Ammate X as a continuous band in the frill with solution of 4 lbs. of Ammate X per gallon of water.


IMPORTANT: Do not apply to desirable plants. Wash sprayer thoroughly after use to remove all Ammate X and to reduce corrosion. Ammate X absorbs moisture readily: keep package tightly closed.

NOTE: 1 lb. is equivalent to 1- $\frac{3}{4}$ cupfuls.

Some species of woody plants are perennial weeds are difficult to control, and retreatment may be necessary if regrowth occurs. Ammate X is water-soluble, non-volatile, and non-flammable. This product is technical material with no additional ingredients.

CONTAINER DISPOSAL: Do not reuse empty container. Destroy it by perforating or crushing and burying in a safe place.

Net. W 2 lbs.



AMMATE X*
NON-SELECTIVE WEED & BRUSH KILLER

The addition of wetting action of dilute sprays

WOODY PLANTS: Bramble and rangelands, a full leaf stage for best results

RATE: Mix 1

TO KILL WEEDS: Locations: and duration conditions.

RATE: Mix 1

ACTIVE INGREDIENTS	
Ammonium Sulfamate	95%
INERT INGREDIENTS	5%

Packaged and Distributed by
STEPHENSON CHEMICAL CO., INC.
COLLEGE PARK, GEORGIA

11-30-71
4267-134

1M 10-71

TO KILL UNDESIRABLE TREES IN NON-CROPPED AREAS. Make a continuous cut or "frill" around the base of the tree with downward axe strokes cutting into the sapwood.

RATE: Apply dry Ammate X to a continuous band at the frill with solution of 1 lbs. of Ammate X per gallon of water.

IMPORTANT: Do not apply to desirable plants. Wash sprayer thoroughly after use to remove all Ammate X and to reduce corrosion. Ammate X absorbs moisture readily; keep package tightly closed.

NOTE: 1 lb. is equivalent to 1- $\frac{3}{4}$ cupfuls.

Some species of woody plants are perennial weeds are difficult to control, and retreatment may be necessary if regrowth occurs. Ammate X is water soluble, non-volatile, and non-flammable. This product is technical material with no additional ingredients.

CONTAINER DISPOSAL: Do not reuse empty container. Destroy it by perforating or crushing and burying in a safe place.

CAUTION:

KEEP OUT OF REACH OF CHILDREN. AVOID CONTACT OF SKIN WITH STRONG SOLUTIONS OF THIS MATERIAL. IN CASE OF CONTACT, WASH OFF WITH PLENTY OF WATER.

EPA Reg. No. 4887-184

Net Wt. 2 lbs.

STEPHENSON

Chemicals

AMMATE X*

NON-SELECTIVE WEED & BRUSH KILLER

CAUTION: KEEP OUT OF REACH OF CHILDREN (READ CAUTIONS ON SIDE PANEL)

*Registered trademark of E. I. Du Pont De Nemours & Company (Inc.)

ACTIVE INGREDIENTS	
Ammonium Sulfamate	95%
INERT INGREDIENTS	5%

The addition of wetting agent or dilute spray

WOODY PLANTS, Bramble and rangelands, a full leaf stage for best results.

RATE: Mix 1


TO KILL WEEDS AT LOCATIONS and duration conditions.

RATE: Mix 1

NOTICE: SEE INSTRUCTIONS ON THE BUYER ASSUMES RESPONSIBILITY WHEN SUCH

Packaged and Distributed by
STEPHENSON CHEMICAL CO., INC.
COLLEGE PARK, GEORGIA

Net. W 2 lbs.



AMMATE X*

**NON-SELECTIVE WEED & BRUSH
KILLER**

DIRECTIONS FOR USE

The addition of a wetting agent, surfactant, or household detergent will improve the wetting action on hard to wet foliage. Use approximately, one teaspoonful to each gallon of dilute spray.

WOODY PLANTS: For control of Poison Ivy, Poison Oak, Poison Sumac, Tree Sprouts, Bramble and other brush in areas, such as, right of ways, roadsides, drainage ditches, rangelands, and pastures—Apply as a drenching spray anytime after the foliage reaches full leaf stage and until foliage begins to discolor due to frost. Wet foliage thoroughly. For best results, spray during periods of high humidity. Respray if any recovery is noted.

RATE: Mix 1 lb. per gallon of water.

TO KILL VEGETATION ON ^(UNSPECIFIED) DRIVEWAYS, WALKS, TENNIS COURTS AND SIMILAR LOCATIONS: Apply uniformly with a sprinkling can or sprayer. The degree of control and duration of effect will vary with weed species, rainfall, temperature and other conditions.

RATE: Mix 1 lb. per gallon of water for 25 sq. ft.

ACTIVE INGREDIENTS	
Ammonium Sulfamate	95%
INERT INGREDIENTS	5%

Packaged and Distributed by
STEPHENSON CHEMICAL CO., INC.
COLLEGE PARK, GEORGIA

Net W 2 lbs

STEPHENSON

Chemicals

AMMATE X*

NON-SELECTIVE WEED & BRUSH
KILLER

**CAUTION: KEEP OUT OF REACH OF
CHILDREN (READ CAUTIONS ON
SIDE PANEL)**

*Registered trademark of E. I. Du Pont De Nemours
& Company(Inc.)

ACTIVE INGREDIENTS

Ammonium Sulfamate

95%

INERT INGREDIENTS

5%

DIRECTIONS FOR USE

The addition of a wetting agent, surfactant, or household detergent will improve the wetting action on hard to wet foliage. Use approximately one teaspoonful to each gallon of dilute spray.

WOODY PLANTS: For control of Poison Ivy, Poison Oak, Poison Sumac, Tree Sprouts, Bramble and other brush in areas, such as, right of ways, roadsides, drainage ditches, rangelands, and pastures. Apply as a drenching spray anytime after the foliage reaches full leaf stage and until foliage begins to discolor due to frost. Wet foliage thoroughly. For best results, spray during periods of high humidity. Respray if any recovery is noted.

RATE: Mix 1 lb. per gallon of water.

TO KILL VEGETATION ON DRIVEWAYS, WALKS, TENNIS COURTS AND SIMILAR LOCATIONS: Apply uniformly with a sprinkling can or sprayer. The degree of control and duration of effect will vary with weed species, rainfall, temperature and other conditions.

RATE: Mix 1 lb. per gallon of water for 25 sq. ft.

NOTICE: SELLER MAKES NO WARRANTY, EXPRESSED OR IMPLIED, CONCERNING THE USE OF THIS PRODUCT OTHER THAN INDICATED ON THE LABEL. BUYER ASSUMES ALL RISK OF USE AND/OR HANDLING OF THIS MATERIAL WHEN SUCH USE AND/OR HANDLING IS CONTRARY TO LABEL INSTRUCTIONS.

Packaged and Distributed by
STEPHENSON CHEMICAL CO., INC.
COLLEGE PARK, GEORGIA

CONTACT OF
MATERIAL IN
WATER.