

Net Weight

This Tomato Dust is effective for control of:

- | | |
|-------------------|------------------------------|
| Tomato Fruit Worm | Septoria Leaf Spot |
| Tomato Pin Worm | Early Blight |
| Tomato Horn Worm | Late Blight |
| Fall Armyworm | Anthracnose |
| | Gray Leaf Spot (Stemphylium) |

DIRECTIONS

Begin applications at first sign of disease. For best results, dust in early morning or late evening when the temperature is low and the air is still. Apply dust to all exposed parts of plants and contact of, especially upper and lower leaf surfaces. Use this container as a shaker with which to apply dust, or better still, use a good dusting machine. Repeat dusting at 5-10 day intervals or as necessary. The measure of control is about 100% for most of plants (about 17 per 100 leaf power).

CAUTION:

Do not apply within 14 days of harvest. Residues must be removed by washing. This product will not control insects if used with lime or other materials which are not compatible.

This product is highly toxic to bees exposed to contact of treated or residue-covered surfaces. Protective information may be obtained from your County Agricultural Extension Service.

Do not use in or near water. Do not use in or near water. Do not use in or near water. Do not use in or near water.

CONTAINER DISPOSAL: Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place.

EPA REG. NO. 188-1



TOMATO DUST

(Contains Sevin and Captan)

CAUTION
DO NOT REACH OF CHILDREN
KEEP OUT OF REACH OF CHILDREN
DO NOT TAKE INTO MOUTH
DO NOT BREATHE DUST

KEEP OUT OF REACH OF CHILDREN AND INSECTS

ACTIVE INGREDIENTS
Sevin (Cyfluthrin) 0.5%
Captan 0.5%
Inert Ingredients 99.0%
Total 100.0%

Manufactured by
STEPHENSON CHEMICAL CO., INC.
COLLEGE PARK, GEORGIA

Manufactured by
STEPHENSON CHEMICAL CO., INC.
COLLEGE PARK, GEORGIA

Net Weight.....

This Tomato Dust is effective for control of:

Tomato Fruit Worm
Tomato Pin Worm
Tomato Horn Worm
Fall Armyworm

Septoria Leaf Spot
Early Blight
Late Blight
Anthracnose
Gray Leaf Spot (Stemphylium)

DIRECTIONS

Begin applications at first sign of disease. For best results, dust in early morning or late evening when the temperature is low and the air is still. Apply dust so all exposed parts of plants are contacted, especially upper and lower leaf surfaces. Use this container as a shaker with which to apply dust, or better still, use a good dust gun. Repeat dusting at 5-10 day intervals or as necessary. The dosage should not exceed 100# per acre of plants (about 1# per 100 foot row).

CAUTION:


Do not apply within 3 days of harvest. Residues must be removed by washing. This product will not control insects if used with lime or other highly alkaline materials.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Co-operative Agricultural Extension Service.

This product is toxic to fish. Keep out of any body of water. Do not contaminate water by cleaning of equipment, or disposal of wastes.

CONTAINER DISPOSAL: Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place.

EPA Reg. No. 4887-153



TOMATO DUST

(Contains Sevin* and Captan)

CONTROLS:

ACTIVE INGREDIENTS:

Captan N[(trichloromethyl) thio]-4-

cyclohexene-1, 2-dicarboximide

Carbaryl (1-naphthyl N-methylcarbamate)

INERT INGREDIENTS:

5.00%

5.00%

90.00%

Total

100.00%

*Trade-Mark of Union Carbide Corp. for Carbaryl (1-naphthyl N-methylcarbamate)

Manufactured by

STEPHENSON CHEMICAL CO., INC.

COLLEGE PARK, GEORGIA

Manufactured by

STEPHENSON CHEMICAL CO., INC.

COLLEGE PARK, GEORGIA

UNLESS THE SPECIAL INSTRUCTIONS
APPEAR, AND IDENTICAL LIST
FOR SEVIN, CAPTAN, AND SEVIN-D
LABEL NO. 4887-153