

WARNING

DO NOT APPLY LINDANE IN POULTRY HOUSES, DAIRY BARNS, OR MILK ROOMS. DO NOT SPRAY OR DIP DAIRY ANIMALS. DO NOT USE ON ANY ANIMAL UNDER THREE MONTHS OF AGE. DO NOT DIP ANIMALS WITHIN 60 DAYS OF SLAUGHTER. DO NOT SPRAY ANIMALS WITHIN 30 DAYS OF SLAUGHTER.

APPLICATIONS FOR YOUNG ANIMALS SHOULD NOT EXCEED 0.03% LINDANE (ONE PINT PER 100 GALLONS OF WATER). DO NOT TREAT SOWS WITHIN TWO WEEKS OF FARROWING OR FOR AT LEAST THREE WEEKS THEREAFTER. AVOID TREATING ANIMALS IN COLD STORMY WEATHER. AVOID TREATING OVER-HEATED OR SICK ANIMALS.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

Highly toxic to fish and wildlife. Do not contaminate any body of water by direct application, cleaning of equipment or disposal of wastes and containers.

WARNING

KEEP OUT OF REACH OF CHILDREN. HARMFUL IF SWALLOWED. AVOID BREATHING VAPOR AND SPRAY MISTS. MAY BE ABSORBED THROUGH SKIN. AVOID CONTACT WITH SKIN. DO NOT LEAVE IN EYES OR ON SKIN. IN CASE OF CONTACT, WASH IMMEDIATELY WITH SOAP AND WATER. AVOID CONTAMINATION OF FEED AND FOODSTUFFS. DO NOT USE ON HOUSEHOLD PETS OR HUMANS.

NOTICE

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

EPA Reg. No. 4887-155

500 10-71

WARNING

DO NOT APPLY LINDANE IN POULTRY HOUSES, DAIRY BARNS, OR MILK ROOMS. DO NOT SPRAY OR DIP DAIRY ANIMALS. DO NOT USE ON ANY ANIMAL UNDER THREE MONTHS OF AGE. DO NOT DIP ANIMALS WITHIN 60 DAYS OF SLAUGHTER. DO NOT SPRAY ANIMALS WITHIN 30 DAYS OF SLAUGHTER.

APPLICATIONS FOR YOUNG ANIMALS SHOULD NOT EXCEED 0.03% LINDANE (ONE PINT PER 100 GALLONS OF WATER). DO NOT TREAT SOWS WITHIN TWO WEEKS OF FARROWING OR FOR AT LEAST THREE WEEKS THEREAFTER. AVOID TREATING ANIMALS IN COLD STORMY WEATHER. AVOID TREATING OVER-HEATED OR SICK ANIMALS.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

Highly toxic to fish and wildlife. Do not contaminate any body of water by direct application, cleaning of equipment or disposal of wastes and containers.

WARNING

KEEP OUT OF REACH OF CHILDREN. HARMFUL IF SWALLOWED. AVOID BREATHING VAPOR AND SPRAY MISTS. MAY BE ABSORBED THROUGH SKIN. AVOID CONTACT WITH SKIN. DO NOT LEAVE IN EYES OR ON SKIN. IN CASE OF CONTACT, WASH IMMEDIATELY WITH SOAP AND WATER. AVOID CONTAMINATION OF FEED AND FOODSTUFFS. DO NOT USE ON HOUSEHOLD PETS OR HUMANS.



100 16, 1971

158, 155

**CAUTION: Do not store below 40° F. for extended period
NET CONTENTS.....**



20% LINDANE

EMULSIFIABLE CONCENTRATE

Contains 1.6 lbs. of Lindane per gal.
Controls FLIES in LIVESTOCK BARNs. TICKS, HORN FLIES,
LICE on ANIMALS. TICKS and CHIGGERS on LAWNS and FIELDS
and certain insects on ORNAMENTALS.

**WARNING: KEEP OUT OF REACH OF CHILDREN
(SEE WARNINGS ON SIDE PANEL)**

ACTIVE INGREDIENTS

Gamma Isomer of benzene hexachloride (from lindane)	20.0%
Xylene	77.0%
INERT INGREDIENTS	3.0%

100.0%

Manufactured by
STEPHENSON CHEMICAL CO., INC.
COLLEGE PARK GEORGIA

Handwritten signature or mark

Handwritten signature or mark

DIRECTIONS

SEE WARNING STATEMENT BEFORE USING

APPLY THIS PRODUCT ONLY AS SPECIFIED ON THIS LABEL.

MIXING DIRECTIONS: When using Lindane Emulsifiable Concentrate, add it to water while agitating the mixture. To prevent the emulsion from settling it should be agitated from time to time. Do not store the emulsion in uncoated steel or iron for periods longer than a day.

FLY CONTROL in LIVESTOCK BARNs — Dilute 1 pt. Lindane 20% to 10 gals. water or 1 gal. Lindane 20% to 80 gals. water. Mix well and spray walls, beams, floors, and all places where insects gather. This treatment will also control roaches, ants and mosquitos if places where the insects hide are treated. Wet all surfaces thoroughly to the point of run-off. Remove animals from barns and cover all feed and water troughs before spraying buildings. Do not spray dairy animals, dairy barns, or milk rooms.

SPRAY or DIP for ANIMALS — Follow dilution rates indicated. Spray dip or treat animals thoroughly. See WARNING statement. Controls the following:

TICKS and HORNFLIES on HORSES and CATTLE—1 tablespoon Lindane 20% to 3 gals. water.

1 pint Lindane 20% to 100 gals. water

LICE on CATTLE and HOGS 1 tablespoon Lindane 20% to 3 gals. water

LICE and TICKS on SHEEP and GOATS . . . 1 pint Lindane 20% to 100 gals. water

SARCOPTIC MANGE on CATTLE and HOGS . . . 2 tablespoons Lindane 20% to 3 gals. water


PSOROPTIC MANGE (SCABIES) on SHEEP . . . 1 quart to 100 gals. water

TICK and CHIGGER CONTROL on LAWNS and FIELDS—Dilute 1 qt. Lindane 20% to 100 gals water. Spray to cover grass, weeds, shrubs thoroughly 1(gallon per 125 sq. ft.) Chiggers frequent areas of grass under trees and near water, especially where brambles are present.

Do not apply to food crops.

Birds feeding on treated areas may be killed.

ORNAMENTALS—Dilute 1 tablespoon Lindane 20% to 3 gals. water or 1 pt. to 100 gals. water. Controls aphids, thrips, leaf miners, and plant bugs.



Nov 16 1971
48-1155

DIRECTIONS

SEE WARNING STATEMENT BEFORE USING

APPLY THIS PRODUCT ONLY AS SPECIFIED ON THIS LABEL.

MIXING DIRECTIONS: When using Lindane Emulsifiable Concentrate, add it to water while agitating the mixture. To prevent the emulsion from settling it should be agitated from time to time. Do not store the emulsion in uncoated steel or iron for periods longer than a day.

FLY CONTROL in LIVESTOCK BARNs — Dilute 1 qt. Lindane 20% to 10 gals. water or 1 gal. Lindane 20% to 80 gals. water. Mix well and spray walls, beams, floors, and all places where insects gather. This treatment will also control roaches, ants and mosquitos in places where the insects hide are treated. Wet all surfaces thoroughly to the point of run-off. Remove animals from barns and cover all feed and water troughs before spraying buildings. Do not spray dairy animals, dairy barns, or milk rooms.

SPRAY or DIP for ANIMALS — Follow dilution rates indicated. Spray dip or treat animals thoroughly. See WARNING statement. Controls the following:

TICKS and HORNFLIES on HORSES and CATTLE— 1 tablespoon Lindane 20% to 3 gals. water.

1 pint Lindane 20% to 100 gals. water

LICE on CATTLE and HOGS 1 tablespoon Lindane 20% to 3 gals. water

LICE and TICKS on SHEEP and GOATS . . . 1 pint Lindane 20% to 100 gals. water

SARCOPTIC MANGE on CATTLE and HOGS . . . 2 tablespoons Lindane 20% to 3 gals. water

PSOROPTIC MANGE (SCABIES) on SHEEP . . . 1 quart to 100 gals. water

TICK and CHIGGER CONTROL on LAWNS and FIELDS—Dilute 1 qt. Lindane 20% to 100 gals. water. Spray to cover grass, weeds, shrubs thoroughly (10 gallon per 125 sq. ft.) Chiggers frequent areas of grass under trees and near water, especially where brambles are present. Do not apply to food crops.

Birds feeding on treated areas may be killed.

ORNAMENTALS— Dilute 1 tablespoon Lindane 20% to 3 gals. water or 1 pt. to 100 gals. water. Controls aphids, thrips, leaf miners, and plant bugs

NOTE: Consult your State Agricultural Extension Service (or County Agent) for additional information, consistent with label directions and cautions.

CONTAINER DISPOSAL: Do not reuse empty container. Destroy by perforating or crushing and burying in a safe place.

