RESTRACTED TEST PLEATERING

For application only by or under the line to the restaurant vision of a certified applicator.

The first tenth of the b

MAR 1 9 1984

The control of the potential resistered gater LPA Reg. No.





20% LINDANE EMULSIFIABLE CONCENTRATE

ONLY FOR SALE TO, USE AND STORAGE BY PEST CONTROL OPERATORS OR OTHER COMMERCIAL AND AGRICULTURAL USERS.

To Control Wood Infesting Beetles of The Families Lyctidae, Anobiidae, and Cerambycidae in Existing Structures.

Also controls FLIES in LIVESTOCK BARNS, TICKS, HORN FLIES, LICE on ANIMALS. TICKS and CHIGGERS on LAWNS and FIELDS and certain insects on ORNAMENTALS.

ACTIVE INGREDIENTS

Contains 16 lbs of Lindane per gal.

KEEP OUT OF REACH OF CHILDREN

WARNING


STATEMENT OF PRACTICAL TREATMENT
TREATMENT CALL A PHYSICIAN OR POISON CONTROL CENTER
MMED ATELY

- I swallowed Contains petroleum distillates. Do not induce vomiting
- If inhared -- Remove victim to fresh air. Apply respiration, if indicated
- to in skin -- Wash area immediately with soap and warm water
- tion eyes Flush with plenty of water and get prompt medical attention SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Reg No 4887-155

EPA Est No 4887-GA-1

Net Contents _____



PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS WARNING

MAY BE ABSORBED THROUGH SKIN, HARMFUL IF SWALLOWED AVOID INHALATION OF SPRAY MIST. AVOID SKIN CONTACT. IN CASE OF CONTACT, WASH IMMEDIATELY WITH SOAP AND WATER. WASH THOROUGHLY AFTER USING AND BEFORE EATING OR SMOKING AVOID CONTAMINATION OF FEED AND FOCD-STUFFS. DO NOT USE ON HOUSEHOLD PETS AND HUMANS.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Birds and other wildlife feeding in treated areas may be killed. Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or-disposal of wastes.

BENEFICIAL INSECT CAUTION—This pesticide is highly toxic to bees exposed to direct treatment or to residues remaining on the treated area. Do not apply when bees are actively visiting the crop, cover crop, or weeds blooming in the treatment area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

PHYSICAL AND CHEMICAL HAZARDS FLAMMABLE. KEEP AWAY FROM HEAT AND OPEN FLAME.

PROTECTIVE CLOTHING REQUIREMENTS

Applicators must wear the following protective clothing during the application process: a light-weight protective suit or coveralls; water-resistant hat; unlined, waterproof gloves; and unlined, lightweight boots. Mixers and loaders aust also wear goggles or a face shield, waterproof gloves and a waterproof apron.

Applicators working in enclosed areas, such as crawl spaces, must wear a respirator approved by OSHA (29 CFR 1910.134).

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

MIXING DIRECTIONS: When using Lindane Emulsifiable Concentrate, add it to water while agitating the mixture. To prevent the emulsion from settling it should be agitated from time to time. Do not store the emulsion in uncoated steel or iron for periods longer than a day.

WOOD INFESTING BEETLES of the Families Lyctidae, Anobildae, and Cerambycidae in homes, barns and other existing structures.

Use a 0.5% emulsion prepared by diluting 3.2 ounces of 20% Lindane to one (1) gallon with water, or 16 ounces (1) mit to 5 gallons of water. Brush solution onto small areas. On large areas apply a course spray until surfaces are wet. Do not apply course sprays in living areas of homes or other existing structures. Do not apply as a mist. Best results are obtained when applied to dry wood. Use in the home is limited to professional licensed. Pest Control Operators only.

Do not use in poultry houses. Do not use in dairy barns and milk rooms. Remove animals from barns and cover all feed and water troughs before spraying buildings. Do not use in edible products areas of food processing plants, restaurants, or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed.

OTHER AGRICULTURAL AND COMMERCIAL USES: Do not apply Lindane in poultry houses, dairy barns, or milk rooms. Do not spray or dip dairy animals. Do not use on any animal under three months of age. Do not dip animals within 60 days of slaughter. Do not spray animals within 30 days of slaughter.

Applications for young animals should not exceed 0.03% Lindane (one pint per 100 gallons of water). Do not treat sows within two weeks of farrowing or for at least three weeks thereafter. Avoid treating animals in cold stormy weather. Avoid treating over-heated or sick animals.

FLY CONTROL in LIVESTOCK BARNS—Difute 1 pt Lindane 20% to 10 gals, water or 1 gal. Lindane 20% to 80 gals, water. Mix well and spray walls, beams, floors, and all places where insects gather. This treatment will also control roaches, ants and mosquitoes if places where the insects hide are treated. Apply a coarse spray until all surfaces are wet. Remove animals from barns and cover all feed and water troughs before spraying buildings. Do not spray dairy animals, Cairy barns, or milk rooms.

SPRAY or DIP for ANIMALS—Follow dilution rates indicated Spray dip or treat animals thoroughly.

Controls the following TICKS and HORNFLIES on HORSES and CATTLE—1 tablespoon Lindane 20% to 3 gals water or 1 pint Lindane 20% to 100 gals water

LICE on CATTLE and HOGS, LICE and TICKS on SHEEP and GOATS 1 tablespoon Lindane 20% to 3 gals, water or 1 pint Lindane 20% to 100 gals, water. SARCOPTIC MANGE on CATTLE and HOGS, PSOROPTIC MANGE (SCABIES) on SHEEP—2 tablespoons Lindane 20% to 3 gals, water or 1 quart to 100 gals, water

TICK and CHIGGER CONTROL on LAWNS and FIELDS—Dilute 1 of Lindane 20% to 100 gals, water Spray to cover grass, weeds, shrubs thoroughly (1 gallon per 125 sq. ft.) Chiggers frequent areas of grass under trees and near water especially where trambles are present. Do not apply to food criss.

GRNAMENTALS—Dilute 1 tablespoon Lindane 20% to 3 gals, water or 1 pt. to 100 gals, water Controls aphids, thrips, real miners, and plant bugs.

NOTE: Consult your State Agricultural Extension Service for County Agents for addition, timformation, consistent with laber directions and cautions.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Do not store below 40°F, for extended period store away from direct sunight in a controller on. Store away from food and feed, Keep container 1 Jht y closed when not in use. Spills or leaking containers should be handled according to Federal. State or tocal procedures under the Resource Contervation and Recovery Act.

PESTICIDE DISPOSAL: Pesticide wastes are toxic improper disposal of excess pesticide, spray mixture, or rinsate is a victation of Federal Law if these wastes cannot be disposed of by use according to label instructions contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance

CONTAINER DISPOSAL: Triple rinse (or equivalent) Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Used dip solutions must be disposed of in accordance with the Resource Conservation and Recovery Act (RCRA). If the applicator generates more than 1000 kg of used dip solution per month or more than 1000 kg used dip solution in combination with other hazardous waste, the material must be treated as a hazardous waste subject to subpart C of RCRA. Any user who wishes to treat, store or dispose of hazardous waste must obtain a permit to serve as a hazardous waste facility tursuant to RCRA.

NOTICE: Seller makes no wurranty, expressed or implied concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and or handling of this material when such use and or handling is contrary to label instructions.



UNITED STATES ENVIRONMENTAL FROTECTION AC NOY

WASHINGTON D.C. 21463

7/ // // / / · ·

1~1.

OFFICE OF PETT BOTT AND TOXIC SUBSTANCES

Mr. W. E. Blasingam Stephenson Chemical Co., Inc. P.O. Box 87188 College Park, GA 30337 " : 111 1 10 1

Dear Mr. Blasingame:

Subject: Lindane RPAR

Stephenson Chemicals 20% Lindane Emulsifiable Concentrate

EPA Reg. No.: 4887-155

Appl. Date: December 23, 1983

The amendment referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable provided that you make the labeling changes listed below before you release the product for shipment bearing the amended labeling:

1. The RPAR Notice incorrectly indicated the language to be used for the RESTRICTED USE PESTICIDE statement. Delete "For application only by or under the direct supervision of a certified applicator" and insert the following at the top of the front panel above the product name. (Note: the type size requirements for the heading "RESTRICTED USE PESTICIDE" are the same as required for the front panel signal word.)

"RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification."

Submit five (5) copies of your revised final printed labeling before you release the product for shipment with the amended labeling.

A stamped copy of the label is enclosed for your records.

If you hold a Federal registration, a supplemental registration, or a Special Local Needs registration, you may continue to produce new stocks of such products, using current labeling, until April 1, 1984. After that date, all remaining inventories of affected products will fall under the Existing Stocks provisions stated in our letter of October 28, 1983.

Sincerely yours,

George I. LaRocca Product Manager (15) Insecticide-Rodenticide Branch Registration Division (TS-767d)