

Net Wt. \_\_\_\_\_



# RoZol REDI-MIX MEAL

## CAUTION

Keep away from humans, domestic animals and pets. If swallowed or absorbed by humans, domestic animals or pets, this material may reduce the clotting ability of the blood and cause bleeding. In that case, intravenous and oral administration of Vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin. Call a physician immediately.

Baits should be placed in areas inaccessible to children, pets, wildlife and domestic animals or in tamper proof bait boxes.

RECEIVED  
SEP 12 973

1571a WI

<b>ACTIVE INGREDIENT:</b>	
2-[ (p-chlorophenyl) phenylacetyl ] -1,	
3-indandione*	0.005%
<b>INERT INGREDIENTS</b>	<b>99.995%</b>
	<hr/>
	<b>100.00%</b>
*Chlorophacinone-Lichadione-Patented in the USA	

Manufactured by  
**STEPHENSON CHEMICAL CO., INC.**  
COLLEGE PARK GEORGIA

100% PURE ROZOL BAIT MEAL

Net Wt. \_\_\_\_\_



# RoZol REDI-MIX MEAL

pets. If  
s or pets,  
blood and  
administra-  
ns are in-  
verdoses of

dren, pets,  
it boxes.

<b>ACTIVE INGREDIENT:</b>	
2-[ (p-chlorophenyl) phenylacetyl ] -1, 3-indandione*	0.005%
<b>INERT INGREDIENTS</b> .....	99.995%
	<hr/>
	100.00%

\*Chlorophacinone-Lichadione-Patented in the USA

Manufactured by  
**STEPHENSON CHEMICAL CO., INC.**  
COLLEGE PARK GEORGIA

4887-158

ACCEPTED

## **CAUTION**

Keep away from humans, domestic animals and pets. If swallowed or absorbed by humans, domestic animals or pets, this material may reduce the clotting ability of the blood and cause bleeding. In that case, intravenous and oral administration of Vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin. Call a physician immediately.

Baits should be placed in areas inaccessible to children, pets, wildlife and domestic animals or in tamper proof bait boxes.

Net Wt. \_\_\_\_\_

**STEPHENSON**

**RoZol REDI-MIX MEAL**

**GENERAL DIRECTIONS**

Place baits in locations where rats and mice feed, water, and/or travel. Bait placements along walls, in corners, and in concealed locations where rats and mice can feed unobserved are desirable. Contaminated or spoiled baits should be replaced promptly.

Protect bait against access of children, pets, livestock and wildlife.

**FOR RATS:** Each station should contain at least 4 ounces of bait. Maintain constant supply of bait at stations for at least 10 days.

**FOR MICE:** Tablespoon amounts ( $\frac{1}{4}$  to  $\frac{1}{2}$  ounce) of bait should be placed at 8 to 12 foot intervals. An uninterrupted supply of bait should be maintained for at least 15 days.

**KEEP AWAY FROM FOODSTUFFS.**

**ACTIVE INGREDIENT:**

2-(p-chlorophenyl) phenylacetyl -1, 3-indandione*	0.005%
<b>INERT INGREDIENTS</b>	<b>99.995%</b>

100.00%

\*Chlorophacinone-Linhadione-Patented in the USA

Manufactured by

**STEPHENSON CHEMICAL CO., INC.**

COLLEGE PARK, GEORGIA