

DIRECTIONS FOR USE (Cont'd)

APPLICATIONS OF THIS PRODUCT IN THE FOOD AREAS OF FOOD HANDLING ESTABLISHMENTS, OTHER THAN AS A SOAK AND CREVICE TREATMENT, ARE NOT PERMITTED.

Spraying unloading docks and outside areas where insects congregate will aid in preventing re-infestations. Sanitation and proper disposal of wastes is essential in any effective insect control program.

PEST CONTROL IN ANIMAL RESEARCH FACILITIES

The use of CARBAMATE 1.5 EC in animal research quarters for pest control will not interfere with the normal activity of the animals.

PEST CONTROL - OUTDOORS

For use by Pest Control Operators and Commercial Applicators for residual control of insects around industrial buildings, homes, garbage areas, dumps, and for mosquito control in non-crop areas. RESIDUAL SPRAY for Ants, Cockroaches, Earwigs, Fleas, Hornets, Millipedes, Mosquitoes, Spiders, Crickets, Clover Mites, Fleas, Scorpions, Sowbugs, Brown Dog Ticks, Wasps and Water-bugs. Mix 8 fluid ounces (1/4 pint) of CARBAMATE 1.5 EC in 1 gallon of water* (1.1% concentration) and apply as a residual spray or with a paint brush to surfaces of buildings, porches, screens, window frames, patios and garages where these insects congregate. For control of wasps and hornets, apply after dark when all of the insects have returned to the nest. Repeat as necessary. Do not treat plants, lawns, flowering plants, shrubs, trees, etc. with this concentration. To control sand flies or punkies (Gnats spp.), mix 8 fluid ounces (1/4 pint) CARBAMATE 1.5 EC in 1 quart of water* (4.5% concentration) and apply with a paint brush to outside surfaces of door and window screens and other screened areas where insects may enter buildings. Thorough coverage is necessary for maximum control.

GROUND MIST SPRAY FOR MOSQUITOES: Apply 4 1/2 to 6 fluid ounces (0.05 to 0.07 lb active) per acre. For mist blower machines calibrated to deliver 100 gallons per hour traveling at a speed of 4 m.p.h. and covering a swath width of up to 350 feet, use 5 1/2 to 8 gallons of spray concentrate per 100 gallons of water* to treat 170 acres. CAUTION: Do not allow to drift on crop land or pasture lands.

AIRCRAFT APPLICATIONS FOR MOSQUITOES: Use CARBAMATE 1.5 EC at dosages of 4 1/2 to 6 fluid ounces (0.05 to 0.07 lb active) per acre in 2 to 4 quarts of water for application in conventional aircraft spray equipment. CAUTION: Do not allow to drift on crop land or pasture lands. NOTE: Due to the high volatility of the solvents in CARBAMATE 1.5 EC, it is necessary to add the formulation directly to the aircraft spray tank and mix well with water. Do not pre-mix in nurse tanks. Aircraft tanks must be closed after mixing to prevent solvent evaporation which would result in crystallization of the active ingredient. Prepared spray should not be held in tank longer than 2 to 3 hours.

PEST CONTROL ON LAWNS AND TURF: Sod Webworms (larva moth): To control sod webworms on lawns and turf, mix 11 fluid ounces (approximately 1/3 pint) of CARBAMATE 1.5 EC in 10 to 15 gallons of water* and apply to 1,000 square feet of area. For best results, mow the lawn and rake dead grass from damaged spots. Water the lawn thoroughly and allow grass to dry before spraying with CARBAMATE 1.5 EC. If possible, do not water again for several days. Three applications at monthly intervals may be required for maximum control. In most areas, the first application should be made in May or June. Consult your local State Agricultural Experiment Station or Extension Service for more specific information regarding timing of applications.

CRICKET BUGS AND HUNTING BILLBUGS (FLORIDA ONLY): Apply 16 fluid ounces (1/2 pint) of CARBAMATE 1.5 EC per 1,000 sq. ft. of lawn or turf area. Do not use less than 15 gallons of water* per 1,000 sq. ft. For best results, use 30 to 40 gallons of water* per 1,000 sq. ft. Repeat as necessary. For effective control of cricket bugs, CARBAMATE 1.5 EC should be applied during the spring and summer months or as recommended by your State Agricultural Experiment Station or Extension Service. For best results, lawns should be watered before applying CARBAMATE 1.5 EC.

PRECAUTIONARY NOTE: Birds feeding on treated areas may be killed. Keep children and pets off treated areas until dry. *The use of this material in water at 40° F. or below will result in crystallization in the spray tank. Do not use in cold water. Do not allow spray mixture to stand overnight in the spray tank.

MIXING CHART Fluid Ounces of Carbamate 1.5 EC to Add To

Concentration Required	1 Gallon U.S.	3 Gallons U.S.	15 Gallons U.S.
1.1%	8.3 fl. oz.	24.9 fl. oz.	124.8 fl. oz.
4.5%	3.3 fl. oz.	9.9 fl. oz.	49.5 fl. oz.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this material other than that stated on the label. Buyer assumes all maximum liability for any and all damage to water bodies and all other living organisms.

January 7, 1983

METALDEHYDE METHIOCARB GRANULES 2-1
CENTRAL PANEL

METALDEHYDE METHIOCARB GRANULES 2-1

Contains 2% Metaldehyde and 1% Mesulcarb

FOR DISTRIBUTION AND USE IN CALIFORNIA ONLY

Active ingredients	
Metaldehyde	2.0%
3,5-Dimethyl-4-methylthio-phenyl methyl carbamate	1.0%
Inert ingredients	97.0%
	Total 100.0%

*Mesulcarb is a registered trademark of the parent company of Farbwerke Bayer, GmbH, Leverkusen.

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Administer milk or water freely and induce vomiting, giving one dose (1/2 oz. or 15 ml. of syrup of ipecac). If vomiting does not occur within 30 minutes, administer second dose. If syrup of ipecac is not available, induce vomiting by sticking finger down throat. Repeat until vomit fluid is clear. Immediately get medical attention.

IF ON SKIN: Remove contaminated clothing and wash skin immediately with soap and warm water.

IF IN EYES: Wash eyes with flowing water for at least 15 minutes.

This pesticide may be fatal to children and to all other pets if eaten. Keep children and pets out of treated areas.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Reg. No. 133-05500

EPA Est. 5481 CA 1

AMVAC CHEMICAL CORPORATION

4100 E WASHINGTON BLVD.

LOS ANGELES, CALIF. 90023

NET WEIGHT: 50 POUNDS