

LINDANE 12-W

Gamma Isomer of Benzene Hexachloride
(from Lindane)

12% WETTABLE POWDER INSECTICIDE

FOR
BEEF CATTLE - HOGS - SHEEP

CONTROLS
Lice, Ticks, Sarcoptic Mange

DIRECTIONS FOR USE:

2 lbs. 3 oz. of Lindane 12W in 100 gallons water equals 0.03% mixture.
3½ teaspoonsful Lindane 12W in 1 gallon water equals 0.03% mixture.

BEEF ANIMALS - To control ticks, lice and sarcoptic mange, use 0.06% mixture for dip or spray. Use 0.03% mixture for young animals. READ CAUTION AND FOLLOW DIRECTIONS.

SWINE - To control lice, spray or dip with 0.03% mixture, 3½ teaspoonsful per gallon water. To control mange (sarcoptic) spray or dip with 0.06% mixture, 7 teaspoonsful per gallon water. READ CAUTION AND FOLLOW DIRECTIONS.


SHEEP - GOATS - To control ticks (Ked), spray or dip with 0.03% mixture, 3½ teaspoonsful per gallon water. To control mange (sarcoptic), spray or dip with 0.06% mixture, 7 teaspoonsful per gallon water. READ CAUTION AND FOLLOW DIRECTIONS.

APPLE - Aphids - 2½ lbs. Lindane 12W in 100 gallons water. May be used as late as petal fall at 2½ lb. dosage for control of rosy aphid. Do not use more than 400 gallons of this spray mixture per acre during any growing season. Do not apply within 60 days of harvest.

PEACHES - Tarnished plant bug and certain other catfacing insects, use 2½ lbs. Lindane 12W in 100 gallons water. *Curculio*, use 2½ lbs. in 100 gallons water in petal fall spray. Repeat application two or three times at about weekly intervals as recommended by State Agricultural Experiment Stations. Do not apply within 60 days of harvest. Do not apply more than 400 gallons of spray mixture per application per acre during spray season.



LINDANE 12-W



ACTIVE INGREDIENTS 12%
Gamma Isomer of Benzene Hexachloride (Lindane)
INERT INGREDIENTS 88%
TOTAL 100%

WARNING

KEEP OUT OF REACH OF CHILDREN
SEE PRECAUTION STATEMENT ON SIDE OF PACKAGE

EPA Reg. No. 4931-48

GOOD-LIFE CHEMICALS, INC.
GOOD-LIFE DRIVE P. O. BOX 637
EFFINGHAM, ILLINOIS 62401

WARNING - IMPORTANT

BEEF CATTLE - Do not dip animals within 60 days or spray within 30 days before slaughter. Do not dip calves under 3 months of age.

HOGS, SHEEP & GOATS - Do not dip within 60 days or spray within 30 days of slaughter. Do not treat young animals under 3 months of age. Do not apply to dairy goats. Do not treat sows within 2 weeks of farrowing nor for at least 3 weeks thereafter. Avoid treating animals in cold stormy weather. Avoid treating overheated or sick animals.

WARNING :

Harmful if swallowed. May be absorbed through skin. May cause irritation of skin and eyes. Avoid inhaling dust, vapor, or mist from sprays. Avoid contact with eyes or skin. Avoid contamination of feed and foodstuffs. Wash thoroughly with soap and water after exposure. For eyes, flush with water and get prompt medical attention.

The Federal Food and Drug Administration has set official tolerances for pesticide residues. Crops having residues in excess of these tolerances should not be marketed without removal of such excess.

This product is toxic to fish and wildlife. Keep out of lakes, ponds, and streams. Do not apply to any area nor use for any purpose not specified on the label.

Do not contaminate water by cleaning of equipment, or disposal of wastes.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

NET CONTENTS 1 POUND

CONTROL NO.

4-1M-73