


CAPTAN - MALATHION




WARNING


KEEP OUT OF REACH OF CHILDREN

See caution statement otherside of package.

U.S.D.A. REG. No. 4931-49



MAY 17 1958



CAPTAN - MALATHION



FOR THE CONTROL OF MANY VEGETABLE INSECTS AND FUNGUS DISEASES

D I R E C T I O N S

Apply GOOD-LIFE CAPTAN-MALATHION dust thoroughly to top and bottom of foliage. It is always best to apply dust early in the morning or evening.

BEANS: To control Mexican bean beetle, leaf hopper, leaf beetle, corn earworm, blister beetle, Western black flea beetle, use 16 pounds per acre.

CUCURBITS: To control striped cucumber beetle, 12-spotted beetle, squash vine borer, melon worm, angular leaf spot, downy mildew, Anthracnose, use 16 pounds per acre.
DO NOT APPLY TO BEANS AND CUCURBITS WITHIN 1 DAY OF HARVEST.

LETTUCE: To control 7 spotted leaf hopper, use 32 pounds per acre.
DO NOT APPLY TO LEAF LETTUCE WITHIN 14 DAYS OF HARVEST
NOR TO HEAD LETTUCE WITHIN 14 DAYS OF HARVEST.

EGG PLANT: To control flea beetle, use 26 pounds per acre.
DO NOT APPLY WITHIN 3 DAYS OF HARVEST.

TOMATOES - POTATOES: To control suck fly, flea beetle, sweet potato beetle, Colorado potato beetle, aphids, young stages of tomato horn worm, early and late blight, Septoria leaf spot, Stemphlium, Anthracnose, use 35 pounds per acre.
DO NOT APPLY TO POTATO AND TOMATOES WITHIN 5 DAYS OF HARVEST.

STRAWBERRIES: To control Botrytis fruit rot, leafspot, aphids and spider mites, use 4 pounds per acre.
DO NOT APPLY WITHIN 3 DAYS OF HARVEST.
ALWAYS WASH VEGETABLES OR FRUIT BEFORE EATING.

ROSES: Apply as a dust weekly to thoroughly cover foliage, top and bottom of leaves (40 to 50 lbs. per acre).
To control blackspot, aphids, rose scale, two spotted spider mite, rose leaf hopper, rose leaf beetle, rose slugs, thrips, rose midges, Japanese beetles and certain other leaf eating insects.

NET CONTENTS 1 POUND

morning or late evening when air is still. Always apply when insects are young and before fungus disease shows up. Insects are easier to control when young, and it is always better to prevent fungus disease than to cure it. Regular application at intervals not to exceed two weeks should give excellent control; however, keep in mind that a heavy rain will wash most of the material off the plants; therefore, as soon as plants are dry, dust should be applied.

IMPORTANT: Best and most economical results will be obtained if a pump dust gun is used which gives a uniform coverage and enables you to easily dust the underside of foliage.

Amount Per Acre	Amount Per 100 Sq. Ft.
16 lbs.	2 1/4 tablespoonsful
26 lbs.	3 1/2 tablespoonsful
32 lbs.	4 1/2 tablespoonsful
35 lbs.	5 tablespoonsful
40 lbs.	5 1/2 tablespoonsful
50 lbs.	6-7 tablespoonsful

C A U T I O N

Avoid contact with skin, eyes and clothing. Avoid breathing of dust. May cause irritation of nose, throat and skin. In case of contact, wash with warm water and soap. For eyes, get medical attention. Store away from children, domestic animals, feed and foodstuffs. If ingested, call a doctor at once and give emetic such as mustard in warm water to induce vomiting.

ALWAYS WASH VEGETABLES BEFORE EATING OR COOKING.
This product is toxic to fish. Keep out of any body of water.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

NOTICE: Seller makes no warranty, expressed or implied concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and or handling of this material when such use and or handling is contrary to label instructions.

Manufactured by
GOOD-LIFE CHEMICALS, INC.
GOOD-LIFE DRIVE, P.O. BOX 687
EFFINGHAM ILLINOIS 62401