

RASPBERRIES AND BLACKBERRIES: For control of Red Berry Mites, Anthracnose, Powdery Mildew, and Rust, spray once when new canes are 8 to 12 inches high at early Spring strength. Repeat in Fall for Red Berry Mites.

CURRENTS AND GOOSEBERRIES: For Powdery Mildew control, use Summer Foliage strength before and after blooming. Do not spray on fruit.

SHADE TREES, SHRUBS AND HEDGES: For control of San Jose Scale, Ash Scale, Rose Scale, Olive Scale, Larch Case Bearer, Leaf Blotch, Maple Gall, Boxwood Canker, Powdery Mildew, Anthracnose, and Nectria Canker, apply at dormant strength. On hedges, spray main stems and limbs. Avoid spraying leaves as much as possible. Do not use on broadleaf evergreens, such as Euonymus and Rhododendrons, or allow spray to drift on these susceptible species. May be used on Spruce and Pines.

SOLD FOR INSECTICIDE AND FUNGICIDE PURPOSES-CAUTIONS: Avoid contact with face and hands. If accidentally splashed into eyes, wash at once with pure water. Do not use an oil spray within three weeks following a Lime-Sulphur spray. Do not use on foliage when temperatures exceed 85° F. Do not spray in the heat of the day. Take care spray drift does not contact light painted surfaces as dark stains or spots may result. Rinse all equipment thoroughly with water after use. **DO NOT STORE WHERE PRODUCT WILL FREEZE.**

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.



SOLUTION 320 BF

LIME SULPHUR FUNGICIDE

ACTIVE INGREDIENTS:
Calcium Polysulphide 30.0%
INERT INGREDIENTS: 70.0%
100.00%

WARNING

KEEP OUT OF REACH OF CHILDREN

See Other Warnings on Back of Label.

NET CONTENT 1 QUART

USDA Reg. No. 4931-64

Packaged By

GOOD-LIFE CHEMICALS, INC.

Good-Life Drive P.O. Box 687
EFFINGHAM, ILLINOIS 62401

Amt. of Spray Desired	Dormant	Fall & Early Spring	Summer Foliage
1 gallon	³ / ₄ pint	5 tablespoons	3 teaspoons
3 gallons	2 ¹ / ₂ pints (5 cups)	¹ / ₂ pint (1 cup)	3 tablespoons
50 gallons	5 gallons	1 gallon	1 quart

DIRECTIONS FOR USE: (See Mixing Table for proper dilutions.)

ROSES: Use at early Spring strength in late Fall after leaves drop, and again in Spring before buds swell, for control of San Jose and Rose Scales. Use Summer Foliage strength during growing season at 10 to 14-day intervals for Powdery Mildew, Red Spider, and Rust control. Observe cautions in hot weather.

APPLES, CHERRIES, PLUMS, AND PEARS: Use at dormant strength before buds swell in the Spring for control of San Jose Scale. For Pear Leaf Blister Mite, delay application until buds begin to swell, as these pests over-winter in the buds. For Scab, Blotch and Powdery Mildew control, use at Summer Foliage strength at cluster bud and prebloom stage. Spray nursery stock and non-bearing fruit trees to clean up these same pests.

PEACHES: For control of Peach Leaf Curl, Brown Rot, San Jose Scale, and Twig Borer, spray at dormant strength after leaves drop in the Fall. Repeat the application in the Spring before the buds begin to swell. Make one application at Summer foliage strength at petal fall to aid in control of Scab and Brown Rot. Do not use on Apricots.

GRAPES: For control of Powdery Mildew and Anthracnose, spray once using foliage strength when new shoots are 4 to 6 inches long. Some injury may occur on existing foliage. Use Dusting Sulfur for rest of season at 7 to 10-day intervals.

MUR-36-1971

49-14 4/10/1971