



4% MALATHION DUST

DIRECTIONS FOR USE:

BEEF CATTLE: Lice and horn fly may be controlled by making thorough application of 4% Malathion Dust—Repeat application in 10 days and as needed thereafter.

DAIRY CATTLE: For the control of horn flies, apply 4 tablespoons (2 oz.) of 4% Dust on the poll, neck and back of animals. Repeat application at 10 to 14 day intervals if needed. Do not apply to dairy animals later than 5 hours before milking or during milking. Do not use Malathion Spray or Dip on dairy animals. Do not contaminate feed, food containers and water troughs. Do not apply to calves under 1 month of age.

DOGS AND CATS: Apply 4% Malathion Dust to thoroughly treat animals, working dust down through hair to the skin, to control fleas, ticks, and lice. Repeat application in 10 to 14 days if necessary. To control fleas in animal quarters (kennels, yard or pen) apply 4% Dust at rate of 1 or 2 pounds per 1000 square feet, using a dust gun to insure even distribution.

POULTRY HOUSES: For the control of northern fowl mites, chicken red mites, and poultry lice, apply 1 pound of 4% Malathion Dust per 50 square feet of litter and floor space, and to roosts, nests and adjacent areas. Repeat as necessary.

POULTRY: After treating poultry house apply 4% Dust to birds using shaker can to insure penetration to skin, to control northern fowl mites, poultry lice and chicken red mites.

SWINE: To control lice on swine use 4% Malathion Dust, making a thorough application to the animals. In addition the pens and houses should also be dusted, using a dust gun to insure even application of 2 pounds per 1000 square feet of surface. Repeat application at 10 day intervals as needed.

CAUTION:

Avoid contact with skin, eyes and clothing. Avoid breathing of dust. May cause irritation of nose, throat and skin. In case of contact, wash with warm water and soap. For eyes, get medical attention. Store away from children, domestic animals, feed and food stuffs. If ingested call a doctor at once and give emetic, such as mustard in warm water to induce vomiting.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

ACTIVE INGREDIENTS:

Malathion (O,O-Dimethyl Dithiophosphate or Diethyl mercaptosuccinate) 4.00%

INERT INGREDIENTS:

TOTAL

96.00%
100.00%

U.S.D.A. REG. No. 4931-65

WARNING

KEEP OUT OF REACH OF CHILDREN.

See other warnings on panel.

Manufactured By

GOOD-LIFE CHEMICALS, INC.

1000 PEEBLES RD., P.O. BOX 100

STURGEON, MINN. 55059

CONTENT: 3 POUNDS