



5% SEVIN* BUG KILLER



* U.S. Patent No. 2,903,478 Union Carbide Chemicals Co.

WARNING

KEEP OUT OF REACH OF CHILDREN

See caution statement otherside of package.

U.S.D.A. REG. No. 4931-72

ACTIVE INGREDIENT:

Carbaryl (1-naphthyl N-Methylcarbamate) 5.00%
 INERT INGREDIENT: 95.00%

TOTAL 100.00%

GRASSHOPPERS

Apply 30 pounds on the above crops excepting cucumbers, melons, pumpkins and squash. A 10 to 20 pound rate is suggested for nymphs on small plants or sparse vegetation in wasteland, rangeland, ditchbanks and borders. A 20 to 30 pound dosage should be used when grasshoppers are mature or material is applied to crops requiring greater coverage.

POULTRY INSECT CONTROL**In or around poultry houses:**

For chicken mite, fleas and bed bugs; Apply 1 pound per 40 square feet of litter or floor space, and to roosts and adjacent areas by means of a shaker can, hand duster or power duster. Force dust into cracks and crevices. Repeat as necessary. Do not treat nest litter. Avoid direct contamination of eggs.

PET INSECT CONTROL**On dogs and cats:**

For ticks and fleas dust liberally over the entire animal, avoiding the eyes, and rub thoroughly into the skin. Comb out dead ticks a few hours after treatment. Repeat at weekly intervals if needed. Avoid treatment of kittens less than 4 weeks of age.

In or around dog and cat quarters:

For ticks and fleas dust liberally in and around sleeping quarters, kennels and dog houses by means of a shaker can, hand or power duster. Force dust into cracks and crevices. Repeat as necessary.

PREHARVEST AND GRAZING USE LIMITATIONS:

No post-treatment time limitations on applications to alfalfa, beans, blueberries, carrots, clovers, cowpeas, cucumbers, egg-plants, forage grasses, grapes, melons, pasture, peanuts, peppers, potatoes, pumpkin, sorghums grown for forage, soybeans, summer squash, sweet corn, tomatoes and winter squash.

Allow 1 day between application and harvest of strawberries.

Allow 3 days between application and harvest of broccoli, Brussels sprouts, cabbage, cauliflower, kohlrabi and lettuce.

Allow 7 days between application to poultry and game birds or poultry and game bird premise treatments, and bird slaughter.

CAUTION!

HARMFUL IF SWALLOWED OR INHALED. Do not take internally. Avoid breathing dust or spray mist. Avoid prolonged or repeated contact with skin. Wash thoroughly after handling. Change contaminated clothing daily. Avoid contamination of foodstuffs. Avoid contamination of feed, feeding troughs and watering receptacles. **KEEP OUT OF REACH OF CHILDREN.**

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

NET CONTENTS: 1 POUND

VEGETABLE INSECT CONTROL:

Suggested dosage refer to pounds of 5% Sevin Bug Killer Dust per acre. (For home gardens, dust for full coverage). Where a dosage range is indicated, use lower rate on young plants; higher rate on mature plants. Use sufficient dust to obtain full coverage. Apply when insects or their damage appear. Repeat at 7-10 day intervals or as necessary unless shorter interval is specified below.

BEANS (Green beans, lima beans, snap beans, cowpeas and black-eyed peas).

Use 10 pounds for Mexican bean beetle; 20 pounds for bean leaf beetle, cucumber beetles, flea beetles, Japanese beetle, leafhoppers, velvet bean caterpillar and western bean cutworm. Use 20 to 30 pounds for armyworm, corn earworm, stink bugs and tarnished plant bug.

CUCUMBER, MELONS, PUMPKIN AND SQUASH

Use 10 to 20 pounds for pickleworm and melonworm; 20 pounds for cucumber beetles, flea beetles, leafhoppers, and squash bugs. Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity. Do not use GOOD-LIFE 5% SEVIN* BUG KILLER on watermelons in Florida.

POTATO, TOMATO, EGGPLANT, PEPPER AND STRAWBERRIES

Use 10 to 20 pounds for Colorado potato beetle, flea beetles and leafhoppers. Use 20 to 40 pounds for European corn borer, fall armyworm, lace bugs, tomato fruitworm, tomato hornworm, tarnished plant bugs and stink bugs, meadow spittlebug, leaf roller and strawberry weevil.

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER AND KOHLRABI

Use 10 to 20 pounds for flea beetles and harlequin bug.

Use 20 to 40 pounds for armyworm, cabbage caterpillars, and corn earworm. Where cabbage looper is the principal problem, use an alternative pesticide that is specifically recommended. For control of small cabbage loopers in home gardens (or in commercial plantings after edible portions begin to form) use GOOD-LIFE 5% SEVIN* BUG KILLER in a 5 to 7 day schedule.

TOBACCO INSECT CONTROL**In plant beds:**

For tobacco flea beetle, apply 1 to 2 pounds per 100 square yards when plants appear. Repeat as needed at 10 to 14 day intervals and just before transplanting.

In Fields:

For hornworms and tobacco flea beetle, apply 1 to 25 pounds per acre. Good coverage is essential. Repeat as needed. Use the highest dosage on large plants. Avoid heavy applications just before priming leaves.

If late applications are necessary, treat immediately after priming rather than before priming. Treated fields may be entered immediately after dust has settled.

Manufactured by
GOOD-LIFE CHEMICALS, INC.

GOOD-LIFE DRIVE, P.O. BOX 687
 EFFINGHAM, ILLINOIS 62401