

**DIRECTIONS FOR USE**

**PREPARATION OF SPRAY:** Fill the spray tank with half the required amount of water, then add the recommended amount of GOOD-LIFE 2 LB. LOW VOLATILE 2,4-D WEED KILLER with agitation and continue filling the tank. Use enough water per acre to give uniform coverage. The amount of water required for low volume applications may vary from 5 to 10 gallons per acre. For high volume applications, 100 gallons or more of water will be needed for good coverage. In any case, use the same amount of 2,4-D recommended per acre.

**GENERAL WEED CONTROL:**

**ROADSIDES, VACANT LOTS AND ALONG FENCE ROWS:** Use 2 to 4 quarts of GOOD-LIFE 2 LB. LOW VOLATILE 2,4-D WEED KILLER in 100 gallons of water per acre. For small areas add 4 tablespoonsful of GOOD-LIFE 2 LB. LOW VOLATILE 2,4-D WEED KILLER to sufficient water to cover 1000 square feet. Thoroughly wet all foliage to runoff.


**PASTURES, MEADOWS, AND RANGELANDS:** Apply 2 quarts per acre of GOOD-LIFE 2 LB. LOW VOLATILE 2,4-D WEED KILLER in sufficient water to provide uniform coverage. Treat forage lands when weeds are growing actively. Do not apply to recently seeded pastures until grass is well established. Avoid application to legumes unless some damage can be tolerated. For control of Wild Garlic, Wild Onions and Canada Thistle, apply 4 to 6 quarts per acre of GOOD-LIFE 2 LB. LOW VOLATILE 2,4-D WEED KILLER in sufficient water for good coverage in early spring and repeat annually until these weeds are under control. Do not apply when grass is in boot to milk stage or after heading begins. Do not graze dairy animals on treated areas within 7 days after applications.

**WOODY PLANT CONTROL:** To control 2,4-D susceptible woody plants such as Willow, Alder, Buckbrush, Sumac, and Elderberry in non-cropped or similar waste areas, use 3 to 4 quarts in 100 gallons of water. Wet thoroughly all parts of the plants, including foliage and stems, to point to runoff.

**LAWNS, GOLF COURSES AND ORNAMENTAL TURF:** Apply 1 1/2 to 2 quarts per acre of GOOD-LIFE 2 LB. LOW VOLATILE 2,4-D WEED KILLER in sufficient water to provide uniform coverage. Do not apply to newly seeded lawns until grasses become well established, and avoid using on Bentgrass, St. Augustine grass, Dichondra and Carpetgrass because of possibility of injury to these grasses and clovers. For control of Wild Garlic and Onion, apply 4 to 6 quarts per acre in very early spring and repeat annually until these weeds are under control.

**FOR TREATING SMALL AREAS:** To treat small areas such as lawns, driveways, garage and incinerator areas and similar small areas - Add 4 tablespoons of GOOD-LIFE 2 LB. LOW VOLATILE 2,4-D WEED KILLER to sufficient water to cover 1000 square feet. Thoroughly wet all foliage to point of runoff.

For best results apply to weeds that are young and in a rapid growing condition. Some woody plants and weeds are hard to kill and repeat applications may be necessary.



**2 LB. LOW VOLATILE 2,4-D WEED KILLER**

(2,4-D Low-Volatile Emulsifiable Concentrate)  
Contains 2.0 Lbs. of 2,4-D Acid Equivalent Per Gallon

**ACTIVE INGREDIENT:**  
Isooctyl Ester of 2,4-D Dichlorophenoxyacetic Acid\* ..... 38.4%

**INERT INGREDIENTS** ..... 61.6%

**TOTAL** ..... 100.0%

\*Equivalent to 2,4-Dichlorophenoxyacetic Acid ..... 25.5%

**SPRAY THE LEAVES - - - KILL THE ROOTS**

Avoid spray drift to susceptible plants such as cotton, tomatoes, flowers, grapes, fruit trees and ornamentals. Spray drift may cause severe injury to both growing and dormant plants. Coarse sprays are less sensitive to drift. Vapors from this product may injure 2,4-D sensitive plants growing near the treated area.

**CAUTION:** Harmful if swallowed. Avoid contact with skin, eyes or clothing. Do not contaminate irrigation ditches or water intended for irrigation or domestic purposes. DO NOT reuse empty container. Destroy by perforating or crushing and burying in a safe place away from water supplies. In case of drums, return to drum conditioner or destroy as listed above.

DO NOT use the same spray equipment for applying other materials to plants as injury may result.

DO NOT store near fertilizers, seed, insecticides or fungicides.

Apply this product only as specified on this label.

**NOTE:** When stored at temperatures below freezing it may be necessary to warm up to at least 45° F. and shake thoroughly before using.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

EPA Reg. No. 4931-81

NET CONTENTS

GALLON

**CAUTION**  
Keep out of reach of Children  
See other precautions on side panels

Manufactured By  
**GOOD-LIFE CHEMICALS, INC.**  
GOOD-LIFE DRIVE P. O. BOX 687  
EFFINGHAM, ILLINOIS 62401

**WEED CONTROL IN CROP:** (See table below) **SMALL GRAIN CROPS** (Wheat, Oats, Barley, Rye) Spray when fully tillered or stooled (4 to 8 inches tall) but before head emerges from the boot. Do not use on grains undersown with legumes except where some legume injury can be tolerated. Do not apply from boot to milk stage or in seedling stage. Do not feed forage or graze treated grain fields within 2 weeks after treatment. Do not feed treated straw to livestock.

**CORN: PRE-EMERGENCE** - Best results are obtained when applied 3 to 5 days after planting but before corn emerges. Do not apply to light sandy soils. **POST-EMERGENCE** - Spray from emergence until corn is 10 inches tall. Above 10 inches in height use corn drops directing the spray down to the base of the corn plant. Do not use from tasseling to dough stage. **SORGHUM - (MILO-SORGO) - POST-EMERGENCE** - Spray when plants are 6 to 10 inches tall when secondary roots are well established. Do not apply during flowering or early dough stage. Above 10 inches in height use drop nozzles directing the spray to the base of the plant.

Amount of Good-Life 2 Lb. Low Volatile 2,4-D Weed Killer Per Acre. Read Complete Directions Before Using

CROP (see detailed directions above)	Average Conditions	Gallons of Spray Per Acre
Small Grains	1 to 1 1/2 qts.	5 to 10 gal.
Corn		
Pre-Emergence	2 to 4 qts.	10 to 20 gal.
Post-Emergence	1/4 to 1 pt.	5 to 10 gal.
Grain Sorghums	1 pt.	5 to 10 gal.
Pastures	1 1/2 to 2 qts.	10 to 20 gal.

**GENERAL INFORMATION:**

GOOD-LIFE 2 LB. LOW VOLATILE 2,4-D WEED KILLER is an oil soluble, water emulsifiable concentrated all purpose weed killer. By addition of water it forms a ready-to-use-spray. It is designed to eliminate weeds from crop lands and lawn areas. The following is a list of some weeds controlled by GOOD-LIFE 2 LB. LOW VOLATILE 2,4-D WEED KILLER:

- |                    |                          |
|--------------------|--------------------------|
| Arrowhead          | Mallows (annual, Poppy)  |
| Beggartick         | Morning Glory            |
| Bindweed           | Mustard                  |
| (hedge, field)     | Nettle, Stinging         |
| Bitter Wintercross | Pennycress               |
| Buckhorn           | Pennywort                |
| Burdock            | Pepper grass             |
| Burhead            | Pigweed                  |
| Chicory            | Plantain                 |
| Cocklebur          | Puncture Vine            |
| Corn Spurry        | Ragweed                  |
| Dandelion          | (Common, Giant, Western) |
| Dry Flower         | Shepard's Purse          |
| Fa. Weed           | Sunflower                |
| Flower-of-the-Hour | Thistle (Roadside, Bulb) |
| Hemp               | Velvet Leaf              |
| Henbit             | Wild Cucumber            |
| Jimson Weed        | Wild Radish              |
| Lamb's-Quart       | Wild Parsnip             |
| Kochia             | Winter Cress             |

**CONTROL NO.**

