


SEVIN★ W.P. 50%



KILLS BUGS
on Vegetables, Fruits,
Ornamental Shrubs, Flowers,
Lawns

CAUTION
KEEP OUT OF REACH OF CHILDREN

See Side Panel For Additional Precautions

USDA REG. No. 4931-91

4931-91 SUBJECT
TO ALL OTHERS.

NOV-26-1968
4931-91
NET CONTENTS 1 POUND

**A Spray Powder for Control of Insects on
Vegetables, Fruits, Ornamental Shrubs,
Flowers, Lawns**

ACTIVE INGREDIENT:

Carbaryl (1-naphthyl N-methylcarbamate) 50% by wt.
INERT INGREDIENTS 50% by wt.
*SEVIN is the registered Trademark of Union Carbide Corp. for
Carbaryl Insecticide.

DIRECTIONS FOR USE:

Use 2 level tablespoons per gallon water and thoroughly wet foliage to control insects on apple, pear, peach, apricot, plum and cherry trees. Do not spray until 30 days after bloom.

Use 2 to 4 tablespoons in one gallon water to control insects on strawberries, raspberries, blackberries, asparagus, beans, peas, corn, lettuce, carrots, beets, parsnips, radishes, turnips, cucumbers, melons, squash, pumpkin, tomato, egg plant, pepper, cabbage, broccoli and potatoes.

Use 2 to 4 tablespoons in one gallon water to spray thoroughly on a wide range of ornamental plants, shrubs, and shade trees including rose, gladiolus, zinnia, chrysanthemum, lilac, arborvitae, juniper, hydrangea, azalea, elm, maple, oak, dogwood, birch and pines. SEVIN injures Boston Ivy.

Controls:

apple aphid	Japanese beetle	plant bugs
bagworm	June beetles	puss caterpillars
birch leaf miner	lace bugs	psyllids
blister beetles	leaf hoppers	rose aphid
boxelder bug	leaf roller	rose slug
boxwood leaf miner	mealy bugs	scale insects
elm leaf aphid	mimosa webworm	tent caterpillars
elm leaf beetle	oak leaf miners	thorn bugs
flea beetles	orange tortrix	thrips (exposed)
gypsy moth	periodical cicada	willow leaf beetles

LAWN INSECT CONTROL: Use 1 cup full of SEVIN 50 in 3 1/2 gallons water for each 100 square feet of lawn surface for control of:

ants	fall armyworm	mosquitoes
clutch bugs	fleas	sod webworms (lawn moths)
earwigs	lathoppers	millipeds

Repeat in 2 or 3 weeks if needed. Use pressure type equipment and apply full water volume to insure good penetration of turf. For best results, mow lawn and apply immediately after rain or watering.

CAUTION: For protection of honey bees avoid use, if possible, during periods when honey bees are visiting the crop. When necessary to use during such period, warn beekeepers well in advance to locate hives at a safe distance until one week after application.

Pre-harvest and grazing limitations —

No post-treatment limitations on applications to almonds, beans, blueberries, carrots, corn torage or fodder, cucumbers, eggplants, filberts, grapes, melons, okra, garden peas, peppers, potatoes, pumpkins, summer squash, sweet corn, tomatoes, walnuts and winter squash.

Allow 1 day between spraying and harvest of apples, cherries, peaches, pears, plums, prunes and strawberries.

Allow 3 days between spraying and harvest of apricots, brocc brussels sprouts, cabbage, cauliflower, head lettuce, kohlrabi, garac beets (roots), nectarines, horseradish, parsnips, radishes, rutabagas, salsify (roots) and turnips (roots), asparagus (Do not repeat applications within three days).

Allow 5 days between spraying and harvest of citrus fruits.

Allow 7 days between spraying and harvest of blackberries, boysenberries, dewberries, loganberries and raspberries.

Allow 14 days between spraying and harvest of Chinese cabbage, collards, dandelion, endive (escarole), garden beets (tops), kale, leaf lettuce, mustard greens, parsley, salsify (tops), spinach, Swiss chard and turnips (tops).

IMPORTANT: HARMFUL IF INHALED OR SWALLOWED. DO NOT BREATHE DUST. DO NOT TAKE INTERNALLY. AVOID PROLONGED OR REPEATED BREATHING OF SPRAY. SKIN CONTACT WITH ANY INSECTICIDE MAY BE HARMFUL! AVOID UNNECESSARY SKIN CONTACT. WASH HANDS BEFORE EATING.

Note for Physician: Carbaryl is a moderate, reversible, Cholinesterase inhibitor. Atropine is antidotal. **Avoid contamination of food.**

KEEP OUT OF REACH OF CHILDREN.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and or handling of this material when such use and/or handling is contrary to label instructions.

Manufactured by
GOOD-LIFE CHEMICALS, INC.
GOOD-LIFE DRIVE, P.O. BOX 687
EFFINGHAM, ILLINOIS 62401