


CAUTION
KEEP OUT OF REACH
OF CHILDREN

SEE OTHER CAUTIONS
ON BACK PANEL
U.S.D.A. REG. No. _____

GOOD-LIFE INSIGNIA

SEVIN* W.P. 50%



BUG KILLER

FOR CROPS - FRUIT - ORNAMENTALS - FLOWERS

LAWN

BERRIES

ACTIVE INGREDIENT:

*Carbaryl (1-naphthyl N-methycarbamate).....50.00%

INERT INGREDIENT.....50.00%
TOTAL 100.00%

*Sevin, U.S. Pat. No. 2,993,473

Registered Trademark of Union Carbide Corp. for Carbaryl Insecticide

Net Contents

Packaged by

GOOD-LIFE CHEMICALS, INC.
GOOD-LIFE DRIVE, BOX 627

DATA SHEET

PRODUCT NAME: ADMINISTRATION

GOOD-LIFE SEVIN* 50 W.P.
U.S.D.A. REG. NO.

FIRST NAME:

GOOD-LIFE CHEMICALS, INC.
GOOD-LIFE DRIVE, BOX 687
EFFINGHAM, ILL. 62401

WAREHOUSE LOCATIONS:

#1 - 735 North First St.
E. St. Louis, Ill. 62201

#2 - Good-Life Drive
Effingham, Ill. 62401

PACKAGING:

2s. 6 x 4 lb.
50 lb. bag

ACTIVE INGREDIENT:

*Carbaryl (1-naphthyl N-methylcarbamate)..... 50.00%

INERT INGREDIENTS..... 50.00%

TOTAL 100.00%

*Sevin, U.S. Pat. No. 2,206,477
Reg. Trademark of Union Carbide Corp. for Carbaryl Insecticide.

This material is purchased and repacked "as is" from Union Carbide Corp.,
New York, N. Y.

February 21, 1967

DIRECTIONS FOR USE:

SEVIN 50 WETTABLE POWDER is to be dispersed in water using constant agitation in the spray tank during application. This product may be applied within 1 day of harvest on all food crops listed unless otherwise indicated.

If SEVIN 50 W. is to be applied by airplane, the dosage of Sevin will be the same as specified for various crops on this label. However, the product should be applied as a dust and care must be exercised to avoid drift of the chemical dust to adjoining fields.

OBSERVE INTERVALS BETWEEN LAST APPLICATION AND HARVEST INDICATED BY NUMBER IN () FOLLOWING CROP.

BEANS: Mexican Bean Beetle - 1 lb. in 20 - 100 gal. water per acre.
Leafhoppers, Bean Leaf Beetle, Cucumber Beetles, Western Bean Cutworm, Flea Beetles, Japanese Beetle, Velvet bean Caterpillar - 2 lbs. in 20 - 100 gal. water per acre.
Corn Earworm, Armyworms, Stink Bugs, Tarnished Plant Bug - 2 to 3 lbs. in 20 - 100 gal. water per acre.

CORN: Corn Earworm, Corn Rootworm Adults, European Corn Borer, Fall Armyworm, Flea Beetles, Japanese Beetle, Flea Beetles, Leafhoppers - 2 to 4 lbs. in 10 - 20 gal. water per acre.
For larvae in whorl (budworm damage) and foliage feeders, apply to entire plant. Repeat as necessary. For insects attacking silks and ears, apply at 2 to 3 day intervals starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. Timing and good coverage are absolutely essential for effective control.

TOBACCO (3): GREEN-LEAFED: Hornworms, Budworms, Flea Beetles, June Beetles, Japanese Beetle - 2 to 4 lbs. in 10 - 20 gal. water per acre.
IN PLANT BEDS: Tobacco Flea Beetle - 2 lbs. per 10 gal. water, apply 6 gal. per 100 sq. yds. Green June Beetle Adults - 1 lb. per 100 gal. water as a trench at 10 to 100 gal. per 100 sq. yds. Avoid excessive application; plant injury may result. Allow three days between application and pruning or cutting.

POTATO, TOMATO, PEPPER: Colorado Potato Beetle, Flea Beetles, Leafhoppers - 1 to 2 lbs. in 10 - 20 gal. water per acre.
Tomato Fruitworm, Fall Armyworm, Tomato Hornworm, European Corn Borer, Stink Bugs, Lace Bugs & Lygus Bugs - 2 to 4 lbs. per acre in 10 - 20 gal. water.

ONION: Corn Earworm, Stink Bugs - 2 to 4 lbs. in 10 - 20 gal. water per acre.

5-07-67

Cucumber Beetles, Squash Bug, Flea Beetles, Leafhoppers - 2 lbs. in 10 - 20 gal. water per acre. Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity. Sevin injures watermelons in Florida.

9/3/67

CARROTS, HORSE RADISH, PARSNIPS, PARSNIPES, RUTABAGAS, BEETS, GALSIFY, TURNIPS - Do not apply within 3 days of harvest. If tops are to be used as food, do not apply within 14 days of harvest. CHINESE CABBAGE (14), COLLARDS (14), DANDELION (14), ENDIVE (ESCAROLE) (14), KALE (14), HEAD LETTUCE (3), LEAF LETTUCE (14), MUSTARD GREENS (14), PARSLEY (14), SPINACH (14), SWISS CHARD (14): Flea Beetles, Harlequin Bug, Leafhoppers - 1 to 2 lbs. per acre in 10 - 20 gal. water. Six-spotted Leafhopper (Aster Yellows Vector) - use 2 to 3 lbs. in 10 - 20 gal. water per acre. Armyworms, Cabbage Caterpillars, Corn Earworm, Harnished Plant Bug, Stink Bugs - 2 to 4 lbs. in 10 - 20 gal. water per acre. Where Cabbage Looper is the principal problem, use an alternative pesticide that is specifically recommended.

ASPARAGUS (1): (On Spears): Asparagus Beetle - 2 to 4 lbs. in 10 - 20 gal. water per acre. Do not repeat application within 3 days. Do not apply within 1 day of harvest.

On Ferns and Brush: Asparagus Beetle, Apache Cicada - 1 to 2 lbs. in 10 - 20 gal. water per acre.


CABBAGE (3), BROCCOLI (3), BRUSSEL SPROUTS (3), CAULIFLOWER (3), KOHLRABI (3): Flea Beetle, Harlequin Bug - 1 to 3 lbs. in 10 - 20 gal. water per acre.

Corn Earworm, Armyworms, Cabbage Caterpillars - 2 to 4 lbs. in 10 - 20 gal. water per acre.

Where Cabbage Looper is the problem, use another pesticide. For small loopers after edible portions form, use Sevin every 7 to 10 days.

APPLES (1), PEACH (1): Scabbing etc., Green Apple Aphid, White Apple Leafhopper, Apple Acalyptus - 1 lb. per 100 gal. water. Apple Maggot, Plum Curculio, Rose Apple Aphid, Woolly Apple Aphid, Red-banded and Fruit Tree Leaf Rollers, Eastern Tent Caterpillar, Pear Psylla, Periodical Cicada, Eye-spotted Bud Moth, European Apple Sawfly, Japanese Beetle, Tentiform Leaf Miner, Forster, San Jose, Pearshell and Lecanium Scale, Bagworm, Green Fruitworm, Apple Rust Mite, Pear Leaf Blister Mite, Pear Sawfly, Harnished Plant Bug - 2 lbs. per 100 gal. water.

RED TAIL SPINACH: On early to mid varieties such as Portland, Gruber, Cinthian, McIntosh, Orleans, Rome Beauty, Puritan, Red Delicious, Minner and Yellow Newton - use 1/2 to 1 lb. per 100 gal. water. Apply for full coverage only once between 10 and 25 days after full bloom. On later to mid varieties such as Baldwin, Ben Davis, Duchess, Early McIntosh, Golden Delicious, Lady Apple, Northern Spy, Rome Beauty, Three Star, Steele Red, Burley, Wealthy, Yellow Transparent and York Imperial - use 1 to 2 lbs. per 100 gal. water. Apply for full coverage only between 10 and 25 days after full bloom. Sevin 40 W may cause Golden Delicious to drop and may injure other varieties.



49; . . .

SPRAYS: Grape Leaf Miner, Grape Leaf Hopper, European Fruit Lecanium,
Grape Leaf Skeletonizer - 2 to 4 lbs. in 100 gal. water per acre.
Grape Berry Moth, Japanese Beetle, June Beetles, Red-banded Leaf Roller -
4 lbs. in 200 gal. water per acre.
May be applied on day of harvest.

SILVERWORM (1), DOG HONEY (7), DEWELEW (7), LOANBERRY (7), RASPBERRY (7): Japanese Beetle, Leaf Rollers, Leaf Toppers, Raspberries, Aphid - 1. In 10 - 20 vol. under per 100.

INSECT AIDS (LAWRENCE, BENTON AND STATE COLLEGE ENTOMOLOGICAL DEPT., IOWA CITY, IOWA) - CYNIPIDAE, CHYTRIDIUM, LILIUS, ANACARDIUM, JUNIPER, MORANGHA, LILIA, AMALTA, ORNITHO, LILIUS, LILIUS, LILIUS (ALLS): Sawflies, Elmwood, Oak and Birch Leaf Miner, Tent Caterpillar, Japanese Beetle, Elm Leaf Beetle, etc. Aphids, Elm Beetles, Plant Bugs, Lace bugs, leafhoppers, Scale insects, Elm Aphid and Slug, Tent Caterpillars, Gypsy Moth and Periodical Cicada, Apple Aphid, Elm or Beetles, Elm Beetles, Leafrollers, deeply hatched, Thrips (exposed), Willow Leaf beetles, Thornbugs, Psyllids, Winesap Webworms, Orange Tortrix, Rust Caterpillars - 2 lbs. per 100 gal or 2 lbs. per gallon.

at 100° at 15 to 20 mm. Hg. on IV5.

LAWN INSECT CONTROL: Anas, Minch, nms, C. nms, G. l. mayworm, flies,
Leafhoppers, crickets, etc. (lawn moths) - 1/1

CAUTION: KEEP OUT OF REACH OF CHILDREN AND ANIMALS.

THIS WOP PHYSICIAN: Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidotal.

Toxic to honey bees. The protection of honeybees, avoid unnecessary use during periods when honeybees are visiting crop. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until 1 week after application.

NOTICE: Because conditions of use which are of critical importance but are beyond our control, 1900-111-100000, 1900, and the GUNITE do not make nor do they authorize anyone involved in the distribution of this product to make any warranty, guarantee or representation, expressed or implied, except that it conforms to its physical description and directions on the label. The GUNITE accepts no responsibility for the express understanding that 1900-111-100000, 1900, or anyone involved in the distribution of this product shall not be held responsible for any loss, risk or liability, personal or otherwise, resulting from its handling, misuse or storage.