



## LINDANE 25-W

(25% Lindane Wettable Powder)

### ACTIVE INGREDIENT:

Gamma Isomer of Benzene Hexachloride  
(From Lindane) ..... 25.0%

INERT INGREDIENTS: ..... 75.0%

TOTAL ..... 100.0%

EPA Reg. No. 4931-127

### WARNING

KEEP OUT OF REACH OF CHILDREN.  
See other warnings on side panel.

Manufactured By

**GOOD-LIFE CHEMICALS, INC.**  
GOOD-LIFE DRIVE P. O. BOX 687  
EFFINGHAM, ILLINOIS 62401

NET CONTENTS 1 POUND

**PREPARATION OF SPRAY:** Maximum dosage .06% or 2 lb. per 100 gal. water (3 tsp. to 1 gal.) Minimum dosage .03% or 1 lb. per 100 gal. water (1½ tsp. to 1 gal.)

ANIMALS	INSECTS	DOSAGE	HOW TO USE	RESTRICTIONS
<u>HOGS</u>	Hog Lice	1½ lb. to 100 gal. water	Spray 1 to 2 qts. per animal.	Do not spray within 30 days of slaughter.
<u>SHEEP &amp; GOATS</u>	Sheep Ticks (Keds)	1½ lb. to 100 gal. water	Spray 1 to 2 qts. per animal.	Do not spray within 30 days of slaughter. Do not use on dairy goats.
<u>BEEF CATTLE</u>	Lice & Ticks	1 - 2 lb. to 100 gal. water	Spray 2 to 4 qts. per animal. Repeated applications are required for tick control.	Do not spray within 30 days of slaughter. Do not apply to dairy animals. Treat calves only at .03% level.
<u>HORSES</u>	Ticks	1½ lb. to 100 gal. water	Spray 2 to 4 qts. per animal. Repeated application required.	

**NOTE:** Since the proper timing of livestock treatments vary in different parts of the country, consult your State Agricultural Extension Service for additional information consistent with label directions and cautions.

Avoid treating animals in cold stormy weather. Avoid treating overheated or sick animals. Do not treat animals under 3 months of age.

This product is toxic to fish and wildlife. Keep out of lakes, streams and ponds.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke and fumes. Do not contaminate water by cleaning of equipment or disposal of wastes.

**WARNING:** Harmful vapor and dust. May be fatal if swallowed. May be absorbed through skin. May produce irritation of skin and eyes. Do not breathe dust or spray mist. Avoid contact with eyes and skin. Wash thoroughly. Avoid contamination of feed or foodstuffs. Apply this product only as specified on this label.

**NOTICE:** Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

CROPS	INSECTS & DISEASES	DOSAGE	HOW TO USE	RESTRICTIONS
<u>SQUASH</u>	Aphids, Cucumber Beetles,	2 lb. per 100	Spray plants thoroughly.	Do not apply after blossoms open. Use on melons and squash late in season may cause reduction in quality.
<u>MELONS</u>	Vine Borer, Stink Bug,	gal. water per		
<u>CUCUMBERS</u>	Melon Worm, Squash Bug, Thrips	acre.		
<u>PEACHES</u>	Plum Curculio	4 lb. per 400	Spray foliage thoroughly. Follow with 1 or 2 additional sprays at weekly intervals as recommended locally.	Do not apply within 60 days of harvest.
<u>PLUMS</u>		gal. water per acre.		
<u>APPLES</u>	Wooly Apple Aphid	2 lb. per 100	Spray foliage thoroughly. Follow with 1 or 2 additional sprays at weekly intervals as recommended locally.	Do not apply within 60 days of harvest.
<u>ONIONS</u>	Aphids, Leafhoppers, Lace	2 lb. per 100	Thorough cover spray at first signs of infestation. Repeat at weekly intervals if infestation persists.	Do not use on spring or green onions.
<u>GLADIOLUS</u>	Bugs, Plant Bugs	gal. water per acre.		
<u>NURSERY PLANTS</u>	Japanese Beetles	2 lb. per 100	Thorough cover spray at first signs of infestation. Repeat at weekly intervals as long as infestation persists.	
<u>GLADIOLUS &amp; DAFFODIL BULBS</u>	Wireworms	2 to 3 oz. per 100 lb. bulbs.	Dust dry bulbs at time of planting.	

ACCEPTED

UNDER THE FEDERAL