



FRONT PANEL


GOOD-LIFE(R)

GARDEN SPRAY OR DUST

ACCEPTED
with COMMENTS
to EPA Letter Dated:

MAR 18 1983

4931-128



ACTIVE INGREDIENTS:

*Methoxychlor, Technical	4.50%
Rotenone	2.00%
Other Cube Resins	3.20%
Zineb (Zinc ethylene bisdithiocarbamate)	6.00%
INERT INGREDIENTS	84.30%
TOTAL	100.00%

*Equivalent to 3.96% 2,2-bis (p-methoxyphenyl)-1,1,1-trichloroethane and 0.54% other isomers and related compounds.

KEEP OUT OF REACH OF CHILDREN

W A R N I N G

See side panels for additional
precautionary statements.

EPA REG. NO. 4931-128

EPA LST. NO. 4931-IL-1

LEFT SIDE PANEL

PRECAUTIONARY STATEMENTSHazards to Humans & Domestic Animals:

WARNING: Harmful if swallowed, inhaled or absorbed through skin. Do not take internally. Avoid breathing dust. Avoid contact with skin. May cause irritation of nose, throat and skin. In case of contact, wash with warm water and soap. Wear clean clothing. For eyes, get medical attention. If ingested, call a doctor at once and give emetic such as mustard in warm water to induce vomiting. Use protective clothing (long pants, long sleeve shirts, gloves and boots) during mixing and loading.

Environmental Hazards

This product is toxic to fish. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from areas treated. Do not use treated forage for feed to dairy animals or animals being finished for slaughter. This product is toxic to bees and should not be applied when bees are actively visiting the area.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE & DISPOSAL

Do not store near food or feedstuffs. Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke and fumes. Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies.

DIRECTIONS FOR USE:

DUST APPLICATION: Apply an even light coating of dust to entire plant. Force it through the foliage so it reaches both sides of leaves. Apply dust when air is still.

SPRAY APPLICATION: Mix amount given with enough water to form a paste. Stir until mixture is smooth and creamy, then dilute with water and pour into sprayer. Mix thoroughly and agitate while spraying to avoid settling of material. Use diluted mixture with good pressure to insure thorough coverage of top and underside of foliage.

AMOUNTS TO USE-DUST APPLICATION- 16 level tablespoonfuls per 200 sq. ft. On strawberries and peas, use one half this amount.

SPRAY APPLICATION 4 oz. to 1 gal. water per 200 sq. ft.

LEFT SIDE PANEL CONTINUED & RIGHT SIDE BEGINS WHEREVER APPROPRIATE

Beans	Bean Aphid, bean leaf beetle, Mexican bean beetle, Thrips.	Anthracnose, Rust, Downy mildew.	DO NOT DUST OR SPRAY WITHIN 7 DAYS OF HARVEST.
Cucumbers, Melons, Squash	Striped cucumber beetle, Vine borer, Pickle worms, Flea beetle, Aphids.	Anthracnose, Alternaria blight, Downy mildew, Blossom blight.	DO NOT DUST OR SPRAY WITHIN 7 DAYS OF HARVEST.
Carrots	Leaf Hopper.	Early and late blights.	DO NOT DUST OR SPRAY WITHIN 14 DAYS OF HARVEST.
Corn (Sweet)	Corn ear worm, Flea beetle, Corn leaf aphid.	Leaf blight, Puccinia rust.	DO NOT DUST OR SPRAY WITHIN 7 DAYS OF HARVEST.
Eggplant	Blister beetle, Flea beetle, Aphids, Eggplant lacebug.	Anthracnose, Fruit rot, Downy mildew, Late blight.	DO NOT DUST OR SPRAY WITHIN 7 DAYS OF HARVEST.
Strawberries	Strawberry Weevil, Leaf Roller.	Leaf Spot, Stem-end Fruit Rot.	DO NOT DUST OR SPRAY WITHIN 14 DAYS OF HARVEST.
Lettuce	Cabbage looper, Six-spotted Leafhopper, Aphids.	Botrytis Slime, Downy mildew.	DO NOT DUST OR SPRAY WITHIN 14 DAYS OF HARVEST.
Peas	Pea Weevil, Aphids.	Anthracnose, Downy mildew, Rust.	DO NOT DUST OR SPRAY WITHIN 10 DAYS OF HARVEST.
Spinach	Aphids.	White Rust.	DO NOT DUST OR SPRAY WITHIN 14 DAYS OF HARVEST.
Tomatoes	Flea Beetle, Aphids.	Early Blight, Late Blight.	DO NOT DUST OR SPRAY WITHIN 7 DAYS OF HARVEST.

ALWAYS WASH VEGETABLES THOROUGHLY BEFORE EATING.

IMPORTANT: Apply early morning or late evening when air is still. Always apply when insects are young and before fungus disease if noted. Regular application at intervals not to exceed two weeks should give excellent control. Repeat application if rain washes dust or spray material off plants. Do not apply when plants are wet.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

4431-185

LEFT SIDE PANEL CONTINUED & RIGHT SIDE BEGINS WHEREVER APPROPRIATE

This container is sold by weight and not by volume. If it does not appear full when opened, it is because contents have settled during shipping.

Manufactured by
GOOD-LIFE, INC.
GOOD-LIFE Drive
P.O. Box 687
Effingham, IL 62401

NET CONTENTS _____

300 91964
17 1

Good-Life, Inc.
One Good-Life Drive
P.O. Box 687
Effingham, IL 62401

Attention: Mr. W. K. Walker

Gentlemen:

Subject: Good-Life Garden Spray or Dust
EPA Registration No. 4931-128
Your Letter Dated February 7, 1983

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, is acceptable subject to the comments listed below. Five copies of the finished labeling must be submitted.

1. In the Environmental Hazard Statement change "Keep... water" to "Do not apply directly to water."
2. Revise the Storage and Disposal Statement to read "Do not reuse empty container (bottle, can, bucket). Wrap (container) and put in trash collection."

A stamped copy of the label is enclosed for your records.

Sincerely yours,

Henry M. Jacoby
Product Manager (21)
Fungicide-Herbicide Branch
Registration Division (TS-767)

Enclosure

TS-767:Jacoby:DCR-03292:WANG-3006C:LM:Raven:479-2013:3/11/83

CONCURRENCES							
SYMBOL							
SURNAME							
DATE							

EPA Form 1320 (4-81) OFFICIAL FILE COPY