

DETERGENT GERMICIDE



Non-Foaming Pipeline Detergent-Germicide
For Dairy C.I.P.

FOR FARM OR INDUSTRIAL USE

ACTIVE INGREDIENTS:

Iodine* 1.6%

INERT INGREDIENTS 98.4%
100.0%

*From Sodium N-Cyclohexyl-N-Palmitoyl Taurate Iodine
Complex and Sodium Dodecyl Benzene Sulfonate-Iodine
Complex

Keep Out of Reach of Children.

DANGER

CORROSIVE. CAUSES EYE DAMAGE AND SKIN IRRITATION.

Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

COMBUSTIBLE: Do not use, pour, spill or store near heat or open flame.

KEEP FROM FREEZING.

EPA Reg. No. 4959-9-AA
EPA Est. No. 4959-IL-01



DIRECTIONS FOR USE:

1. **PRE-RINSE.** Immediately after milking, thoroughly rinse the entire pipeline system with sufficient 110°F. (45°C.) water. Permit water to discharge to drain. Do not re-circulate rinse water. In particularly hard water areas, it is advisable to slightly acidify the rinse water by using one ounce of DYNE to twenty gallons of water.

CLEAN. For most satisfactory results, we recommend the use of DYNE-MATE®. Consult label for proper cleaning use directions.

RINSE. Following cleaning cycle, rinse line thoroughly with lukewarm water. In hard water areas, add one ounce of DYNE to twenty gallons of water.

4. **SANITIZE.** Immediately before milking, sanitize with one ounce of DYNE to ten gallons of lukewarm water, permitting two minutes contact time. (This provides 12½ ppm titratable iodine.) For spray applications use one ounce of DYNE to five gallons of water. (This provides 25 ppm titratable iodine.) Rinse with potable water and drain, if required by state or local regulations. Follow all state and local health department regulations.

DYNE is its own indicator of germicidal activity. The amber color at use dilution shows the presence of the active ingredient — iodine.

Not adversely affected by common types of water hardness or by cold water.

STORAGE AND DISPOSAL

Storage: Keep Container closed when not in use. Do not store below 0°F or above 100°F. for extended periods.

Disposal: Do not contaminate water, food, or feed by storage or disposal, or cleaning of equipment. In case of spill, flood area with large quantity of water, washing to sewer or collection vessel.

Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

Do not reuse empty container. Triple rinse, puncture, and dispose of container in a sanitary landfill or by incineration.

© Copyright, 1988, West Agro, Inc.

0108

NET CONTENT: 15 GALLONS

WESTAGRO®


WEST AGRO, INC. / KANSAS CITY, MO 64153

002-0108-B0888

MADE IN U.S.A.



LOT NO:



Not Reviewed. Registrant claims to be in accordance with
Reregistration of Fungicide Product Guidelines for Phase 2
Response, Page 2-12, Active to Inert Change in Status