

Reg # 4959-15

PM-32

12/1

**STORAGE AND DISPOSAL**

**Storage:** Keep container closed when not in use. Do not store below 25° F or above 100° F for extended periods.

**Dispose:** Do not contaminate water, food or feed by storage or disposal, or cleaning of equipment. In case of spill, flood area with large quantity of water, washing to sewer or collection vessel.

Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

Do not reuse empty container. Triple rinse, puncture, and dispose of container in a sanitary landfill or by incineration.

**Danger:** Causes eye damage and skin irritation. Do not get in eyes, on skin, or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food.

**First aid:** In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Call a physician immediately.

**Note to physician:** Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression, and convulsions may be needed.

SOLVAY Animal Health, Inc.  
Mendota Heights, MN 55120-1139 USA  
Distributed from Charles City, IA 50616-9989 USA  
3-7086-8808  
List No. 7005

# Superdyne® II

Disinfectant

For farm and industrial use only

For cleaning, disinfecting, and sanitizing animal housing and equipment; for sanitizing poultry drinking water; for use as a shoe bath prior to entering barns and poultry houses; for use as a deodorant; for sanitizing food-processing equipment



**First aid:** Flush eyes thoroughly with water. If swallowed, drink plenty of water. Do not induce vomiting.

**Precautions:** Avoid contact with eyes, skin, and clothing. Do not breathe vapors.

**Warning:** 1.7% available iodine

**Keep out of reach of children**

**Danger:** Causes eye damage and skin irritation. Harmful if swallowed.

**See also panel for first aid and additional precautions.**

**Net Content: 1 Gallon (128 fl. oz./3.785 liters)**

SOLVAY Animal Health, Inc.

SPECIMEN

BEST AVAILABLE COPY

Superdyne II Disinfectant is tuberculocidal at a dilution of 3 ounces (90 ml) of Superdyne II Disinfectant to 5 gallons (19 liters) of water.

Superdyne II Disinfectant is not adversely affected by water hardness or low-temperature water.

Superdyne II Disinfectant has its own built-in indicator of germicidal activity. (When the amber color disappears, a fresh solution should be prepared.)

#### Directions for use

**Sanitizing buildings and equipment on the farm:** To sanitize surfaces of buildings, equipment, or utensils which have previously been thoroughly cleaned, use 1 ounce (30 ml) of Superdyne II Disinfectant in 5 gallons (19 liters) of water. Spray or swab solution on surfaces to be sanitized. Allow to drain dry without rinsing.

**Sanitizing poultry drinking water:** Add 1 ounce (30 ml) of Superdyne II Disinfectant to every 10 gallons (38 liters) of drinking water. The regular use of the solution keeps the water clean and free of bacteria and other harmful organisms.

**Flood spray disinfection:** Use 1 ounce (30 ml) of Superdyne II Disinfectant in each 5 gallons (19 liters) of water as the final sanitizing step on previously cleaned and processed equipment and utensils. Do not use for flood spray disinfection of buildings. Superdyne II Disinfectant does not require a final rinse. **U.S. PATENT OFFICE REGISTRATION NO. 178,187**

**Shoe bath:** To help prevent tracking disease organisms into poultry houses, farrowing houses, and hog barns, place a shoe bath inside the doorway, containing 3 ounces (90 ml) of Superdyne II Disinfectant per gallon (3.8 liters) of water. After scraping shoes outside doorway, stand in shoe bath for 30 seconds prior to entering building interior. Change shoe bath daily.

**Deodorant:** Superdyne II Disinfectant destroys many odors as it sanitizes and disinfects. For general deodorant applications (garbage pails, refuse containers, etc.), swab or spray on a solution of 1 ounce (30 ml) of disinfectant per gallon (3.8 liters) of water.

**Cleaning and disinfecting buildings and equipment on the farm:** Before proceeding as indicated below, remove all animals and feeds from premises, cars, boats, trucks, and other equipment. Remove all litter and manure from floors, walls, and surfaces of barns, pens, stalls, chutes, and other facilities and fixtures occupied or traversed by animals. Empty all troughs, racks, and other feeding and watering appliances, covering all surfaces with Superdyne II Disinfectant solution. Remove all bedding, ropes, and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers. Wash all buildings, cars, boats, and other stored objects. Do not leave livestock or poultry equipment with Superdyne II Disinfectant on it. All stored feed racks, mangers, troughs, automatic feeders, buckets, and containers must be thoroughly rinsed with potable water prior to reuse.

Then clean and disinfect in one easy step with Superdyne II Disinfectant. Use 1/2 ounce (15 ml) of Superdyne II Disinfectant per gallon (3.8 liters) of water. Spray solution on to be cleaned and disinfected with a pump sprayer or with a power sprayer. Start at the highest point of the building or equipment being cleaned and work down. Let solution drain dry without rinsing.

BEST AVAILABLE COPY