

LFI sanitizer

A low-foaming iodophor for sanitizing in the beer-bottling, soft drink-bottling, milk-bottling, and food-canning-industries.

***ACTIVE INGREDIENTS:**

Butoxy polypropoxy polyethoxy ethanol-iodine complex 12.47%
 Polyethoxy polypropoxy polyethoxy ethanol-iodine complex 0.37%
INERT INGREDIENTS: 87.16%
 Total 100%

*(providing 1.6% titratable iodine)

CAUTION: Keep Out of Reach of Children.
 (See additional cautions on side panel.)

NET CONTENTS:

EPA Reg. No. 4959-18

U. S. PATENT NOS. 3,650,965



WEST AGRO-CHEMICAL, INC.
 SHAWNEE MISSION, KANSAS 66222

ACCEPTED
 4959-18
 JUL 16 1981

Under the Federal Insecticide, Fungicide, and Plant Disease Act as amended, for the pesticide registered under EPA Reg. No.

DIRECTIONS FOR USE

Equipment Sanitizing:

Pre-clean equipment with cold or lukewarm water. Fill with solution containing 2 ounces of LFI SANITIZER per 10 gallons of water (provides 25 ppm titratable iodine). When ready for use, drain off sanitizing solution.

Tank and Vat Sanitizing:

Pre-clean with water (not hot). Add a solution containing 2 ounces of LFI SANITIZER per 10 gallons of water (provides 25 ppm titratable iodine). After five minutes, drain off solution.

Sanitization of Walls, Floors, Conveyors, and other similar surfaces:

Clean daily with solution containing 2 ounces of LFI SANITIZER per 10 gallons of water (provides 25 ppm titratable iodine). Clean weekly, or at end of batch run with solution containing 4 ounces of LFI SANITIZER per 10 gallons of water (provides 50 ppm titratable iodine). Rinse.

Sanitizing of Beverage Containers:

To sanitize previously cleaned beverage containers, use a solution containing 1 ounce of LFI SANITIZER per 10 gallons of water (provides 12½ ppm titratable iodine). Invert container, drain, and allow to air-dry.

DIRECTIONS FOR DILUTION

Titratable Iodine

1 oz. LFI SANITIZER per 10 gallons water	=	12½ ppm
2 oz. LFI SANITIZER per 10 gallons water	=	25 ppm
4 oz. LFI SANITIZER per 10 gallons water	=	50 ppm

LFI SANITIZER is regulated under the Food Additives Amendment of the Federal Food, Drug, and Cosmetic Act for use on food-processing equipment and utensils up to 25 ppm of titratable iodine, without requiring a rinse with potable water.

CAUTION: AVOID CONTACT WITH FOOD.

AVOID CONTACT WITH SKIN OR EYES. In case of contact with eyes, flush immediately with water; get medical attention.

Rinse empty container thoroughly with water and discard.

MADE IN U.S.A.

BEST AVAILABLE COPY

PL 1182

4959-18

1871