

FOR INDUSTRIAL USE ONLY

LFI

A low-foaming iodophor for sanitizing in the beer-bottling, soft drink, and food-canning industries.

ACCEPTED

MAY 27 2003

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pest control, disinfecting, and sanitizing uses, and EPA Reg. No. 4959-18

ACTIVE INGREDIENT:

Iodine* 1.6%

INERT INGREDIENTS: 98.4%

Total 100.0%

* From Butoxy polypropoxy polyethoxy ethanol-iodine complex and Polyethoxy polypropoxy polyethoxy ethanol-iodine complexes

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

FIRST AID

If in Eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes. Then continue rinsing. • Call a Poison Control Center or doctor for treatment advice.
If Swallowed	<ul style="list-style-type: none"> • Call a Poison Control Center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the Poison Control Center or doctor. • Do not give anything to an unconscious person.
If on Skin or Clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. Call a Poison Control Center or doctor for treatment advice.
If Inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a Poison Control Center or doctor for treatment advice.
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency treatment information.</p>	
NOTE TO PHYSICIAN	
<p>Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsions may be needed.</p>	

EPA REG. NO.: 4959-18

EPA EST. NO. 4959-IL-01

Sold By: WEST ACRO, INC.
11100 N. Congress Ave.
Kansas City, Mo. 64153

0001-0403

Made in U.S.A.

PRECAUTIONARY STATEMENTS: Hazards to Humans and Domestic Animals

CAUTION: Avoid contact with food. Causes moderate eye injury. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling.

ENVIRONMENTAL HAZARDS:

This product is toxic to fish and aquatic organisms. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other water unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Fresh sanitizing solution should be prepared daily or more often if the solution becomes diluted or soiled.

EQUIPMENT SANITIZING:

To pre-clean equipment, wash with cold or lukewarm water with a solution of 1 ounce LFI SANITIZER to 5 gallons water. Rinse with potable water. To sanitize, spray or immerse equipment with a fresh solution of 1 ounce LFI SANITIZER to 5 gallons cold or lukewarm water (provides 25 ppm titratable iodine). Allow a one minute contact time. When ready to use, drain sanitizing solution.

TANK AND VAT SANITIZING:

Pre-clean with 1 ounce LFI SANITIZER to 5 gallons water (not hot). Rinse with potable water. To sanitize, spray or fill vat with 1 ounce LFI SANITIZER to 5 gallons water (provides 25 ppm titratable iodine). After 5 minutes, drain off solution.

SANITATION OF WALLS, FLOORS, CONVEYORS, AND OTHER SIMILAR SURFACES:

Pre-clean surface by scraping or flushing to remove soil. Then thoroughly clean with a solution 2 ounces of LFI SANITIZER to 5 gallons of water. Rinse with potable water. Sanitize with a solution containing 1 ounce of LFI SANITIZER to 5 gallons of water (provides 25 ppm titratable iodine). Allow 2 minutes contact time. Do not rinse.

SANITIZING OF BEVERAGE CONTAINERS:

To sanitize previously cleaned beverage containers, which were first washed in detergent and rinsed, use a solution containing 1 ounce LFI SANITIZER to 5 gallons of water (provides 25 ppm titratable iodine). Allow for a one minute contact time. Invert containers, drain and allow to air dry.

DIRECTIONS FOR DILUTIONS:

1 oz. LFI SANITIZER to 5 gallons water	=	25	ppm
2 oz. LFI SANITIZER to 5 gallons water	=	50	ppm

STORAGE AND DISPOSAL

STORAGE: Keep container closed when not in use. Do not store below 32 F or above 100 F for extended periods.

DISPOSAL: Do not contaminate water, food, or feed by storage or disposal, or cleaning of equipment. In case of spill, flood area with large quantity of water, washing to sewer or collection vessel. Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

Do not reuse empty container. Triple rinse, puncture and dispose of container in a sanitary landfill or by incineration.

NET CONTENTS:

LOT NO.:

4959-18

5/27/2003

Page 181