



DRAFT LABELING
Submitted 8/18/88 revised from 2/29/88



UDDERSAN®
Sanitizer and Udder Wash

Active Ingredients:

Iodine*	1.75%
Phosphoric Acid	11.50%

Inert Ingredients:	86.75%
--------------------------	--------

Contains Glycerin

100.00%

* From Nonylphenoxypolyethoxyethanol-Iodine complex

Keep Out of Reach of Children
DANGER

Statement of Practical Treatment:

If Swallowed: Drink promptly a large quantity of water. Do not induce vomiting. Avoid alcohol. Get medical attention.

If In Eyes: Flush with plenty of water for 15 minutes. Get medical attention.

If On Skin: Wash with plenty of soap and water. Get medical attention if irritation persists.


Note to Physician: Probably mucosal damage may contraindicate the use of gastric lavage.

Precautionary Statements:

Hazards to humans and domestic animals.
DANGER: Concentrated product is harmful or fatal if swallowed. Avoid contact of undiluted product with skin, eyes and clothing. Undiluted product causes eye and skin damage. Wear goggles and rubber gloves when handling. Harmful or fatal if swallowed. Avoid breathing vapors or fumes. Wash thoroughly with soap and water after handling.

Environmental Hazards:

This pesticide is toxic to fish. Keep out of lakes, ponds, streams or other public waterways without an appropriate NPDES permit. For NPDES permits, contact your regional EPA office, or the appropriate agency in your state. Do not contaminate water by cleansing of equipment or disposal of wastes.



BEST AVAILABLE COPY

DIRECTIONS FOR USE:

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

For Use as an Udder Wash:

- Prepare an Uddersan solution using 1 ounce of sanitizer to each 5 gallons of lukewarm water (100 degrees F.). This solution contains 25 ppm titratable iodine.
- Dip a single service paper towel into wash solution and wash teats clean to the base of the udder.
- Forestrip.
- Dry teat with individual paper towel. Attach milker unit.
- Prepare a fresh solution if wash solution fades noticeably or becomes diluted or soiled.

For Sanitizing Dairy Equipment and Utensils:

- Wash the equipment to be sanitized using SUPER KLEENITER[®] Manual Cleaner, following label instructions. Rinse with potable water and drain.
- Prepare a solution by mixing 1 ounce of UdderSan Sanitizer for each 5 gallons of water, or add 1/2 ounce of sanitizer to a 10 quart pail of water. This provides 25 ppm titratable iodine.
- Immerse equipment--milking claws, liners, teat cups into solution making sure that all surfaces are in contact with the solution. A contact time of one minute is required.
- For equipment that cannot be immersed (bulk coolers, receiver groups, etc.) spray a 25 ppm iodine solution onto all pre-cleaned surfaces for a sparkling, spot-free shine.
- Allow solution to drain and air-dry.
- Prepare a fresh solution daily or more frequently if sanitizing solution fades noticeably or becomes diluted or soiled.

UdderSan Sanitizer and Udder Wash is its own indicator of germicidal activity. The amber or yellow color shows the presence of the active ingredient-- iodine.

BEST AVAILABLE COPY

Draft Labeling EPA Submission
Revised 8/18/88 from 2/29/88
UDDERSAN®

Page 3

STORAGE AND DISPOSAL: Keep container closed when not in use. Do not store below 20 degrees F. or above 100 degrees F. for extended periods. In case of spill, flood areas with large quantities of water. Product or rinsates that cannot be used should be diluted with water before disposal in a sanitary sewer. Do not reuse container, but place in trash collection. Do not contaminate food or feed by storage, disposal or cleaning of equipment.

CSE/j1
8/18/88
GRA.7:Uddersan

BEST AVAILABLE COPY