

**DIRECTIONS FOR USE (continued)**

**FOR FOOD EGG SANITIZING** - Thoroughly clean all eggs. Thoroughly mix 5 oz. of this product with 10 gallons of warm water to produce a 200 ppm available chlorine solution. The sanitizer temperature should not exceed 130°F. Spray the warm sanitizer so that the eggs are thoroughly wetted. Allow the eggs to thoroughly dry before casing or breaking. Do not apply a potable water rinse. The solution should not be re-used to sanitize eggs.

**STORAGE AND DISPOSAL**

**THOROUGHLY RINSE ALL MEASURING DEVICES AND STORE OUT OF REACH OF CHILDREN.**

Store this product in a cool, dry area away from direct sunlight and heat to avoid deterioration. In case of spill, flood area with large quantities of water. Product or rinsates that cannot be use should be diluted with water before disposing in a sanitary sewer. Do not contaminate food or feed by storage, disposal or cleaning of equipment. Thoroughly rinse empty container with water and discard by placing the container in trash collection.

**PRECAUTIONARY STATEMENTS:**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**



**Corrosive.** May cause severe skin and eye irritation or chemical burns to broken skin. Causes eye damage. Wear goggles or safety glasses when handling this product. Wear rubber gloves when handling

this product. Wash after handling. Avoid breathing vapors. Vacate poorly ventilated areas as soon as possible. Do not return to area until strong odors have dissipated.

**ENVIRONMENTAL HAZARDS:**

This pesticide is toxic to fish and aquatic organisms. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or public waters unless this product is specifically identified and addressed in an NPDES permit. Do not discharge effluent containing this product to sewer systems without previously notifying the sewage treatment plant authority. For guidance, contact your State Water Board or Regional Office of the EPA.

**PHYSICAL AND CHEMICAL HAZARDS: STRONG OXIDIZING AGENT.** Mix only with water according to the label directions. Mixing this product with organic matter (e.g. urine, feces, etc.) or chemicals (e.g. ammonia, acids, detergents, etc.) will release chlorine gas which is irritating to eyes, lungs, and mucous membranes.

EPA Reg. No. 4959-40  
EPA Est. No. 4959-IL-01

**FOR INDUSTRIAL USE ONLY**

**WEST AGRO  
CHLORINATING SANITIZER**


**ACTIVE INGREDIENT**

Sodium Hypochlorite ..... 6.6%

**INERT INGREDIENTS** ..... 93.4%

**TOTAL** ..... 100.00%

**KEEP OUT OF REACH OF CHILDREN**



**STATEMENT OF PRACTICAL TREATMENT (First Aid)**

**IF CONTACT WITH EYES OCCURS:** Flush eyes with plenty of water for at least 15 minutes. Get prompt medical attention.

**IF CONTACT WITH SKIN OCCURS:** Wash with plenty of soap and water. Remove and wash contaminated clothing before reuse.

**IF SWALLOWED:** Drink large amounts of water. DO NOT induce vomiting. Call a physician or poison control center immediately.


See additional precautions on side panel.

West Agro, Inc.  
11100 North Congress  
Kansas City, Missouri 64153  
Reorder# 1-800-421-1905

**HAZARD RATING**

- 4-EXTREME
- 3-HIGH
- 2-MODERATE
- 1-SLIGHT
- 0-INSIGNIFICANT

**TOXICITY**



Chemtrec Emergency Number  
1-800-424-9300

Made in USA

**A MEMBER OF THE WORLDWIDE  
Tetra Laval Group**

**DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

**NOTE:** This product degrades with age. Use a chlorine test kit and increase dosage, as necessary, to obtain the required level of available chlorine.

**FOR SANITIZING FOOD PROCESSING OR DAIRY EQUIPMENT:**

**CLEAN-IN-PLACE METHOD** - Thoroughly clean equipment after use. Prepare a volume of a 200 ppm available chlorine solution equal to 110% of volume capacity of the equipment by mixing the product in a ratio of 5 oz. product with 10 gallons of water. Pump solution through the system until full flow is obtained at all extremities, the system is completely filled with the sanitizer and all air is removed for at least 10 minutes to insure contact with all internal surfaces. Remove some cleaning solution from drain valve and test with a chlorine test kit. Repeat entire cleaning/sanitizing process if effluent contains less than 50 ppm available chlorine.

**RINSE METHOD** - A solution of 100 ppm available chlorine may be used in the sanitizing solution if a chlorine test kit is available.

Solutions containing an initial concentration of 100 ppm available chlorine must be tested and adjusted periodically to insure that the available chlorine does not drop below 50 ppm. Prepare a 100 ppm sanitizing solution by thoroughly mixing 2½ oz. of this product with 10 gallons of water. If no test kit is available, prepare a sanitizing solution by thoroughly mixing 5 oz. of this product with 10 gallons of water to produce approximately 200 ppm available chlorine by weight.

Clean equipment surfaces in the normal manner. Prior to use, rinse all surfaces thoroughly with the sanitizing solution, maintaining contact with the sanitizer for at least 2 minutes. If solution contains less than 50 ppm available chlorine, as determined by a suitable test kit, either discard the solution or add sufficient product to reestablish a 200 ppm residual. Do not rinse equipment with water after treatment and do not soak overnight.

Sanitizers used in automated systems may be used for general cleaning by may not be re-used for sanitizing purposes.

**FOR DISINFECTING FARM PREMISES** - Remove all animals, poultry, and feed from premises, vehicles, and enclosures.

Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, chutes and other facilities occupied or transversed by animals or poultry. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. To disinfect, saturate all surfaces with a solution of at least 1000 ppm available chlorine for a period of 10 minutes. A 1000 ppm solution can be made by thoroughly mixing 25 oz. of this product with 10 gallons water. Immerse all halters, ropes and other types of equipment used in handling and restraining animals or poultry, as well as the cleaned forks, shovels and scrapers used for removing litter and manure. Ventilate buildings, cars, boats, and other closed spaces. Do not house livestock or poultry or employ equipment until chlorine has been dissipated. All treated feed racks, mangers, troughs, automatic feeders, fountains and waterers must be rinsed with potable water before reuse.

**NET CONTENTS:  
LOT NUMBER:**

**BEST AVAILABLE COPY**