

300 | 174958
17 | 2

JUN 26 1986

Mr. Dan Briscoe
Administrative Manager
Young's Incorporated
P.O. Box 71
Roaring Spring, PA 16673

Dear Mr. Briscoe:

Subject: Amendment - Label Revisions/Product Chemistry
Young's Long and Lean Rid Fly
EPA Registration No. 4987-6
Your Letter Dated April 18, 1986

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable provided that you make the labeling changes listed below before you release the product for shipment bearing the amended labeling:

- Following the "Directions for Use" heading, insert the following wording, "It is a violation of Federal law to use this product in a manner inconsistent with its labeling."

The following comment pertains to product chemistry.

- Submit a revised Confidential Statement of Formula which includes certified limits on all active ingredients listed on the label. (Note: this does not include listed minerals.)

50575:Shell:LR-10:KEMCO:6/19/86:6/26/86:TAR:VO

SYMBOL ▶							
SURNAME ▶							
DATE ▶							

Submit five (5) copies of your final printed labeling before you release the product for shipment.

A stamped copy of the labeling is enclosed for your records.

Sincerely yours,

George T. LaRocca
Product Manager (15)
Insecticide-Rodenticide Branch
Registration Division (TS-767C)

Enclosure

young's Long & Lean Aid Fly



EPA REGISTRATION NO. 4987-6
 EPA ESTABLISHMENT NO. 4987-PA-01
 (Contains RABON® Oral Larvicide)

To prevent the development of Horn Flies, Face Flies, House Flies and Stable Flies in the manure of treated cattle.

ACTIVE INGREDIENT:

2-Chloro-1-(2, 4, 5-trichlorophenyl)
 vinyl dimethyl phosphate.....

JUN 23 1986

.973%*

INERT INGREDIENTS.....

*Each pound contains 4.47 g RABON.

**Refers only to ingredients which are not larvicidal.

RABON is a registered trademark of Shell Oil Company.

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended for the pesticide registered under EPA Reg. No. 4987-6

50 LBS. NET — FEEDING DIRECTIONS ON BACK

GUARANTEED ANALYSIS

MINERALS

CALCIUM, not more than	14.00%
CALCIUM, not less than	12.00%
PHOSPHORUS, not less than	10.00%
MAGNESIUM, not less than	5.00%
SULPHUR, not less than	1.00%
POTASSIUM, not less than	.90%
SALT, not more than	16.50%
SALT, not less than	15.50%
IRON, not less than	.23%
ZINC, not less than	.32%
MANGANESE, not less than	.19%
COPPER, not less than	.057%
COBALT, not less than	.0057%
IODINE, not less than	.010%
SELENIUM, not less than	.0004%

VITAMINS

VITAMIN A (U.S.P. Units/lb.)	150,000
VITAMIN D (U.S.P. Units/lb.)	75,000
VITAMIN E (I. Units/lb.)	100

INGREDIENTS

Monocalcium Phosphate, Dicalcium Phosphate, Bone Phosphate, Calcium Sulfate, Magnesium Oxide, Calcium Carbonate, Salt, Ethylene Diamine Dihydrochloride (source of Iodine), Cobalt Sulfate, Zinc Oxide, Zinc Sulfate, Copper Sulfate, Iron Sulfate, Manganous Oxide, Manganese Sulfate, Vitamin A Acetate, D-Activated Animal Sterol (source of Vitamin D), dl-Alpha Tocopheryl Acetate (source of Vitamin E), Dried Cane Molasses, Yeast Culture, Sodium Selenite, Iron Oxide (coloring agent), Mineral Oil.

WARNING: Do not permit intake of supplemental Selenium to exceed 0.1 PPM on a complete ration basis.

KEEP OUT OF THE REACH OF CHILDREN

CAUTION

Read the entire label for use directions and other precautionary statements.

**PRECAUTIONARY STATEMENTS
 HAZARDS TO HUMANS**

CAUTION: Keep out of the reach of children. Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water. If irritation persists see a physician.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not add directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in a dry place in original container.

Container Disposal: Completely empty bag into mixing equipment. Then dispose of empty bag in a sanitary landfill or by incineration or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.

Manufactured By **young's, inc.** P.O. Box 71, Paring Spring, PA 16673

DIRECTIONS FOR USE
young's LONG & LEAN RID FLY

CATTLE: (For Mixing In Feed)

Feed Mineral Mix at the rate of 1.87 ounces per 750 pounds of body weight per head per day to beef and dairy animals by mixing it into the ground grain or complete dairy ration.

— OR —

CATTLE: (For Self-Feeding)

Allow free choice to cattle. 1200 pound cattle should consume an average of 3 ounces of this product per day. This will supply the recommended average daily intake of 70 mg. of RABON per hundred pounds of body weight. Animals should not be starved for minerals before initiation of this Mineral Mix product but OMIT all salt from other feed after the program is started. Do not feed other free-choice sources of salt or mineral containing salt in any form. Do not feed RABON in other feeds when using this self-feeding (free-choice) method. Allow one mineral feeder for each 15-20 head, and place near watering and loafing areas. Put out no more than one month's supply at a time, and protect from rain. If consumption of LONG & LEAN RID FLY is above 3 ounces per day, reduce the number of feeding locations or relocate. If consumption is below 3 ounces per day, increase the number of feeding locations.

This mineral mix containing RABON may be fed up to slaughter and to lactating dairy cows. Start feeding this product early in the spring before flies begin to appear and continue feeding throughout the summer into the fall until cold weather restricts fly activity.

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Rations containing this product may be fed up to slaughter and to lactating dairy cows without withholding the milk from market during or after treatment.

Start feeding RABON larvicidal feeds early in the spring before flies begin to appear and continue feeding throughout the summer and into the fall until cold weather restricts fly activity.

When fed, this product passes through the digestive system into the animal's manure where it kills fly larvae on contact shortly after fly eggs hatch.

It prevents the development of horn flies, face flies, house flies, and stable flies in the manure of treated cattle and house flies in treated swine, but is not effective against existing adult flies.

In some cases, supplemental fly control measures may be needed in and around cattle lots and barns to control adult house flies and stable flies which can breed not only in manure but in other decaying vegetable matter or silage on the premises.

In order to achieve optimum fly control, this product should be used in conjunction with other good management and sanitation practices.

This product will mix uniformly in feeds when standard mixing procedures are followed. Thus, usual problems that are common to all feed preparation and which cause stratification, such as excessive free-fall or excessive handling, are to be avoided. It is recommended that appropriate preblending techniques be employed to assure adequate distribution throughout the feed mix. The premix should be preblended with ground grain, protein supplements, mineral mixes, etc., before being added to roughages, such as chopped hay or silage. Mixing time should be adequate to assure uniform dispersion. Optimum performance can be assured only if this product is dispersed uniformly with the feedstuff at the recommended level. Common feed mixing equipment (i.e., vertical mixers, horizontal blenders, mixer/feeder truck) may be used to prepare formulated feeds.

Feeds prepared with this product should not be pelleted, nor be mixed with feeds containing predominantly pellets. Further, this product should not be mixed in liquid feed supplements.

