

(A list of ...
(all of ...
(or other ...

(...
(...

Panel

(...

ACCEPTED
MAY 12 1967
UNDER THE FEDERAL INSECTICIDE
AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 011-75 SUBJECT
TO ATTACHED COMMENTS.

ACCEPTED
5-12-67
5011-75
UNDER THE FEDERAL INSECTICIDE
CONTROL ACT
REGISTERED
SUBJECT
TO ATTACHED COMMENTS.

all be the equivalent of a 1% DDT (Finisect products).

Application Method and Rate: To stop total area to be treated.

Apply 5 ml of diluted General Formula M-15 per acre (0.05 lb. actual DDT)

to kill adult mosquitoes and gnats. Pay particular attention to all zones

of stagnation ponds, etc. Standing water should be treated for mosquito

larvae with a suitable larvicide. IMPORTANT: Shrimp and crab may be

killed at application rates recommended on this label.

areas such as stables, barns, rubbish piles, etc. Do not use in dairy barns
or on livestock intended for slaughter.

Note: Use of this product as for outdoors and in stables and barns should be
limited to pest control operators or trained personnel.

Most successful fogging is done after sunset, in the evening, when the earth
has to a temperature less than that of the overlying air. Disperse fog in
the fog by using the ground and drifting widely through grass and foliage.

Use prevailing wind direction to plan fogging operation. Start at back end
of area to be treated, and work into the wind at cross angle. If wind becomes
strong enough to be directly against the fog, directions of operation should be
reversed.

When fogging in a stable, use the fogging machine in the center of the
stable, moving from end to end, etc. Do not use in dairy barns
or in areas where livestock are kept.

Note: Use of this product as for outdoors and in stables and barns should
be limited to pest control operators or trained personnel.



For killing flies in stables and barns (other than dairy), use one pint of

diluted Formula MU-15 to treat 8,000 cubic feet. The air movement in the stable or barn should be reduced by closing doors, windows, and other openings. Avoid contaminating water, feed or food stuffs, covering same with wrapping paper if necessary. Humans should vacate and animals removed from premises before indoor treatment with the fog and not re-enter until premises have been ventilated. Do not fog in areas animals have received direct applications of DDVP formulations within 8 hours.

Keep all unprotected persons out of operational area. Poisonous by skin contact, inhalation, or swallowing. Rapidly absorbed through skin. If on skin, wash immediately with soap and water. Do not get in eyes, on skin, or on clothing. In case of contact, immediately remove all contaminated clothing and flush skin or eyes with plenty of water for at least 15 minutes. For eyes, get medical attention. Do not breathe fog or spray mist. Wash thoroughly with soap after handling and before eating or smoking. Wear clean clothing. Atropine is antidotal. Keep away from heat and open flame. Do not treat in areas where food or feed crops are growing.

Antidotes: If swallowed, give a tablespoon of salt in a glass of warm water and repeat until vomit fluid is clear. Have victim lie down and keep quiet. Call a physician immediately.

This product will kill fish. Keep out of lakes, streams, and ponds.



5-12-67
5011-75
UNDER THE FEDERAL INSECTICIDE ACT
FOR THE REGISTRATION OF PESTICIDES
EPA REG. NO. 105-100000-01

WARNING SYMPTOMS:

Symptoms include weakness, headache, tightness in chest, blurred vision, non reactive pin-point pupils, salivation, sweating, nausea, vomiting, diarrhea, and abdominal cramps.

NOTE TO PHYSICIAN

Treatment: Atropine is the specific therapeutic antagonist of choice

parasympathetic stimulation, Atropine Sulfate should be injected at 10 minute intervals, in doses of 1 to 2 milligrams, until complete atropinization has occurred. Morphine is contraindicated. Clear chest by postural drainage. Oxygen administration may be necessary. Observe patient continuously for 48 hours. Repeated exposure to Cholinestrase inhibitors may, without warnings, cause prolonged susceptibility to very small doses of any cholinestrase inhibitor. Allow no further exposure until time for cholinestrase regeneration has been allowed as determined by blood test.

NON-WARRANTY

Seller makes no warranty of any kind, expressed or implied, regarding the use of this product. The buyer agrees that the seller shall not be responsible or liable for injury to crop, soil, person, or property resulting out of the use or handling of the product, whether or not in accordance with the label directions or not, and accepts and uses the product at his own risk and conditions.

IMPORTANT STORAGE INSTRUCTIONS

Keep in original container in a cool, dry place, away from sunlight and moisture.
Store in a cool, dry place, away from sunlight and moisture. Do not use after the expiration date.
Storage until reaching room temperature and then use immediately.

Manufactured by:

P O Box 406
Westfield, Indiana 46074

Net Contents: 35 pounds

ACCEPTED
5-2-67
SP-11-75
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POLICE
ED. UNO...
COMMENTS: