

[5041-1]

INSECTICIDE

KWIK-KILL

— KILLS —

ROACHES, WATER BUGS, MOTHS, FLIES,
MOSQUITOES, ANTS, SILVERFISH, BEDBUGS,
CARPET BEETLES, BUFFALO MOTHS,
and certain other insects

ACTIVE INGREDIENTS 100%

Contains: Deodorized Petroleum Distillate, Technical
Piperonyl Butoxide,* Pyrethrins

* (Butylcarbityl) (6-propylpiperonyl) ether and related
compounds

USDA Reg. No. 5041-1

Manufactured by
SUBURBAN PEST CONTROL CO.
HYATTSVILLE, MARYLAND
WARfield 7-2307

CAUTION

Keep out of reach of Children

DIRECTIONS

ROACHES, WATERBUGS, ANTS AND SILVERFISH

Spray KWIK KILL so as to wet all cracks, crevices, and under tables, behind sinks, drainboards and cabinets; or wherever these insects may be seen or known to be. This will drive out and kill insects contacted.

FLIES AND MOSQUITOES—

For use as a space spray close all windows and doors. Spray KWIK KILL upwards to produce a thick mist. After about 10 minutes the dead insects may be swept up and the windows and doors opened.

CLOTHES MOTHS—

Closets, chests, drawers, etc., should be thoroughly sprayed before clothing is placed in them. Garments should be carefully sprayed both inside and out, paying particular attention to the folds and seams. In spraying delicate fabrics the sprayer nozzle should be held at least 2 feet from the material being treated. Rugs should be sprayed sufficiently on both sides so as to wet the surface. Articles not stored in tight containers should be re-treated at monthly intervals.

CARPET BEETLES AND BUFFALO MOTHS—

In rooms infested with these insects in addition to the procedure used for clothes moths, particular attention must be paid to direct the spray into all cracks in the flooring and behind baseboards.

BEDBUGS

Carefully treat all bedspreads and mattresses, paying special attention to all hiding places. In the case of heavy infestations it may be necessary to spray all cracks and crevices about the room.

REPEAT APPLICATIONS AS NEEDED

CAUTION — DO NOT SPRAY INTO AN OPEN FLAME

Remove or cover fish bowls and aquaria before and for approximately one hour after treatment.

Harmful if swallowed — Keep away from children