AQUACIDE

pellets

A WEED KILLER TO KILL OFFSHORE WEEDS IN RECREATIONAL LAKES AND PONDS



ACCE Represent of the reach of children

EPA Reg. # 5080-2

NOV 3 0 1982

enjoy fishing!

EPA Est. # 5080-WI-1

Under the Federal Insert

Pengicide, and Rodan SEE OTHER PRECAUTIONS
or amounted, for the proON SIDE PANEL

directed rada Con SID

EPA Reg. No. 5080-2

ACTIVE INGREDIENTS:
Sodium, 2,4-Dichlorophenoxyacetate*
INERT INGREDIENTS:
Equivalent to 17.5% 2,4-Dichlorophenoxyacetic acid

100%

CONTENTS 10 POUNDS

AQUACIDE CO. 801 First Street Hudson, WI 54016



AQUACIDE IS INTENDED TO TREAT SUBMERGED WEEDS IN RECREATIONAL LAKES AND PONDS.

GENERAL DISCUSSION OF SUBMERGED WEED CONTROL:

Many types of weeds are found in lakes and ponds. In the formulation of *AQUACIDE* the herbicide 2.4-D is predominantly for control of broad leafed plants.

AQUACIDE is intended to control submerged perennial water weeds, such as Water Thyme, Dutch Moss, Milfoil, Fanwort, Tape Grass, Naiad, Ludwiga and certain other pond weeds.

Probably the most important attribute of AQUACIDE is that the pelletform stratifies the herbicide at the bottom of the lake. This enables weed control while the weeds are very small and just appearing from the bottom.

The stratification does limit the weed killing power in the fall inasmuch as most top sprays are more effective with blooming weeds.

AQUACIDE will not kill algae or top floating weeds. The effectiveness of AQUACIDE is reduced by mud and silt. It will function best in a firm bottomed area.

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

The AQUACIDE pellets are hard and dissolve slowly in water. The proper way to spread the herbicide pellets is to broadcast them with a fanning motion of the hand over the surface of the water, similar to the way grass seed is sown.

Apply AQUACIDE in the spring and early summer when new growth is just beginning. Effects will be visible in 7 to 10 days and full kill will take place in 3 to 5 weeks. Occasionally a second treatment will be necessary in areas of

heavy infestation and should be made in 3 to 5 weeks, or when weeds show signs of recovering. Control will last from 1 to 3 years.

Sow 2½ pounds per 1000 square feet of surface in areas of average weed density. Applications of 4 lbs. per 1000 square feet may be used in heavily infested areas. Do not contaminate water used for spraying, irrigation or potable purposes.

PRECAUTIONARY STATEMENTS CAUTION

HAZARDS TO HUMANS (and Domestic Animals): Avoid contact with skin, eyes and clothing. Wear gloves when spreading pellets.

ENVIRONMENTAL HAZARDS: Treatment of dense weed areas can result in oxygen depletion due to decomposition of dead weeds. This oxygen loss can cause fish suffocation. To avoid fish suffocation when treating an entire body of water, treat 1/3 to 1/2 of the area at a time and allow 3 to 5 weeks before continuing to untreated areas. Water drawn directly from treated areas may damage desirable plants. Do not contaminate water by cleaning of equipment or disposal of wastes. Consult State Fish and Game Agencies before applying this product to public waters.

storage and disposed: Store unused pellets in a dry location. Pesticide or rinse water that cannot be used according to label directions must be disposed of according to applicable Federal or State procedures. Triple rinse empty container and offer for recycling or puncture and dispose of in a sanitary landfill, or by other approved State and local procedures.

AQUACIDE CO. 801 First Street Hudson, WI 54016