

Horning's Bacteriostatic Fabric Softener is intended for use in commercial laundries only.

Fabric treated as directed will develop residual bacteriostatic activity against Gram positive (*S. aureus*) organisms under conditions of high humidity or wet contamination. Horning's Bacteriostatic Fabric Softener is not effective against Gram negative organisms.

Horning's Bacteriostatic Fabric Softener inhibits the growth of ammonia producing bacteria (*B. ammonia-ferens*) which act on urine in the diapers to produce ammonia.

Horning's Bacteriostatic Fabric Softener softens cloths, thus eliminating chafing, "Static Cling" out of nylons and other synthetics. Improves ironing time.

Rinse empty container thoroughly with water and discard it.



BACTERIOSTATIC FABRIC SOFTNER

FABRIC SOFTNER CONCENTRATE
RESIDUAL BACTERIOSTAT UNDER
CONDITIONS OF HIGH RELATIVE
HUMIDITY OR WET CONTAMINATION.

ACTIVE INGREDIENTS:

n-Alkyl (60% C₁₄, 30% C₁₆, 5% C₁₂, 5% C₁₈)
dimethyl benzyl ammonium chlorides 5.0%

n-Alkyl (68% C₁₂, 32% C₁₄) dimethyl
ethylbenzyl ammonium chlorides 5.0%

INERT INGREDIENTS: 90.0%
100.0%

Weight: 100 lb. Drum EPA Reg. No. 5145-16

ACCEPTED
SEP 28 1974 See right side panel for additional precautions
and First Aid.
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 5145-16

DANGER

KEEP OUT OF REACH OF CHILDREN

and First Aid.

HORNING'S SAFETY PRODUCTS CO.
1180 W. 28th Street
INDIANAPOLIS, INDIANA

USE DIRECTIONS:

Use 1½ ounces Horning's Bacteriostatic Fabric Softener per 100 pounds of laundry in RINSE CYCLE.

DANGER:

Keep Out of Reach of Children. Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food.

FIRST AID:

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN:

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

000000

ACCEPTED

SEP 24 1957

UNDER THE PROVISIONS OF THE
LAND REVENUE AND DISTRICT ACT
OF 1956 (1956) POLSON REGISTER
SERIAL NO.