

DE VERE

QD QUATERNARY DISINFECTANT

DISINFECTANT

DEODORIZER

SANITIZER

ACTIVE INGREDIENTS:

A kyl (50% C14, 40% C12,
10% C16) dimethyl benzyl ammonium chlorides 10.0%

INERT INGREDIENTS: 90.0%

TOTAL INGREDIENTS: 100.0%

DANGER:

Keep Out of Reach of Children. Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling concentrate. Harmful if swallowed.

Avoid contamination of food.

See antidote statement on left panel.

DE VERE CHEMICAL CO., INC.

JANESVILLE, WISCONSIN 53545

1923 BELOIT AVENUE

ACCEPTED

4-6-73

5174-18

COLD DISINFECTION

For disinfection of previously cleaned surgical instruments, barber tools and dental equipment, simply submerge instruments for a minimum time of ten minutes in a solution containing 1½ ounces QD per gallon of water.

RESTAURANT AND BAR RINSE

- 1. Scrape and prewash utensils and glasses whenever possible.**
- 2. Wash with DE VERE LIME SOLVENT.**
- 3. Rinse with clean water.**
- 4. Sanitize in a solution of one ounce DE VERE QD to four gallons of water (200 ppm). Immerse all utensils for at least two minutes or for contact time specified by governing sanitary code. No Terminal Potable Water Rinse Required.**
- 5. Place sanitized utensils on a rack or drainboard to air dry.**

FIRST AID:


In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.


If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately. Rinse empty container thoroughly with water and discard.

EPA REG. NO. 5174-18

CONTENTS: 1 Gallon

USE DIRECTIONS:





USE DIRECTIONS:

HOUSEHOLD

For sanitizing and disinfecting floors, walls and inanimate hard surfaces. For Schools, Homes, Locker Rooms, Garbage Pails, Sink Tops, Corridors, Classrooms, Offices and Shower Stalls. Apply solution with mop or cloth. Sanitize with 1 ounce to 4 gallons of water. Disinfect with 1 ounce to 2 gallons of water.

HOSPITAL

For floors, walls and other hard surfaces. In Nursing Homes, Institutions, as well as Sink Tops, Garbage Pails, Telephones and Restrooms, use 1 ounce per 1 gallon of water. At this level (800 ppm active quat.) QD will be effective against *Pseudomonas aeruginosa*. All surfaces must be thoroughly wet with the recommended use dilution by wiping, mopping or scrubbing the area.

FOOD PROCESSING EQUIPMENT

Clean and rinse articles Prior to sanitizing application. For sanitization of precleaned food processing equipment and food utensils, dilute 1 ounce per 4 gallons of water to provide 200 ppm of active quaternary. At this level, No Potable Water Rinse is required.

DAIRIES

To sanitize dairy equipment, first clean and rinse the equipment thoroughly. Then apply sanitizing solution containing 1 ounce to 4 gallons of water (200 ppm). At this level, No Potable Water Rinse is required. Follow recommendations of local Health Board.

ACCEPTED WITH COMMENTS