

*QD Quaternary Disinfectant*

### DIRECTIONS FOR USE

**GENERAL PURPOSE SANITIZATION AND DISINFECTION.** For sanitizing and disinfecting floors, walls and other hard non porous surfaces. For schools, homes, locker rooms, garbage pails, sink tops, corridors, classrooms, offices and shower stalls. Apply solution with mop or cloth. Sanitize with 1 ounce QD to 4 gallons of water. Disinfect with 1 ounce QD to 2 gallons of water.

**HOSPITAL DISINFECTION.** For floors, walls and other hard non porous surfaces in nursing homes, institutions, as well as sink tops, garbage pails, telephones and restrooms, use 1 ounce QD to 1 gallon of water. At this level, 800 ppm active quat, QD will be effective against *Pseudomonas aeruginosa*. All surfaces must be thoroughly wet with the recommended use dilution by wiping, mopping or scrubbing the area.

**FOOD PROCESSING EQUIPMENT.** Clean and rinse articles prior to sanitizing application. For sanitization of pre-cleaned food processing equipment and food contact surfaces, use 1 ounce QD to 4 gallons of water to provide 200 ppm of active quaternary. At this level, no potable water rinse is required.

**DAIRIES.** To sanitize dairy equipment, first clean and rinse the equipment thoroughly. Then apply sanitizing solution containing 1 ounce QD to 4 gallons of water at 200 ppm. At this level, no potable water rinse is required. Follow recommendations of the Health Board.

#### GENERAL CLASSIFICATION

This classification is subject to law to use this product in a manner consistent with its labeling.

#### RESTAURANT AND BAR RINSE:

1. Scrape and prewash dishes and glasses whenever possible.
2. Wash with De Vere DIME SOLVENT.
3. Rinse with hot water.
4. Sanitize with 1 ounce De Vere QD to four gallons of water, 200 ppm quaternary, for at least two minutes or for contact time specified by manufacturer's label. For terminal, potable water rinse required.
5. For sanitizing dishes, sink, a tank or drainboard, spray.

# DE VERE

## QD QUATERNARY DISINFECTANT

DISINFECTANT      DEODORIZER      SANITIZER

#### ACTIVE INGREDIENT:

Alkyl (50% C14, 40% C12, 10% C16) dimethyl benzyl ammonium chloride

#### INERT INGREDIENTS:

TOTAL INGREDIENTS:

**KEEP OUT OF REACH OF CHILDREN**

**DANGER**

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

ACCEPTED  
OCT 25  
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the purposes registered under EPA Reg. No. 5174-12

DE VERE CHEMICAL CO., INC.

JANESVILLE, WISCONSIN 53547

1923

ONLY FOR SALE TO USE AND STORAGE  
BY SERVICE PERSONS

**COLD DISINFECTION:**

For disinfection of previously cleaned surgical instruments, barber tools and dental equipment, simply submerge instruments for a minimum time of ten minutes in a solution containing 1 ounce QD per gallon of water.

**STORAGE AND DISPOSAL**

**STORAGE:** Store in a cool, dry place. Store in original container only.

**DISPOSAL:** Do not contaminate water, food, or feed by storage or disposal. Do not use empty container. Triple rinse, or equivalent. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

EPA Reg No. 5174-18

EPA Est. No. 5174-WI-1

NET CONTENTS ONE GALLON

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS AND  
DOMESTIC ANIMALS**

**DANGER: CAUSES EYE DAMAGE  
CAUSES SEVERE SKIN IRRITATION,  
ESPECIALLY TO BROKEN SKIN  
HARMFUL IF SWALLOWED**

Do not get in eyes, on skin or on clothing.  
Wear goggles or faceshield and rubber gloves when handling this product.  
Remove and wash contaminated clothing.  
Do not take internally.

**STATEMENT OF PRACTICAL TREATMENT**

**IF IN EYES:** Flush with plenty of water for at least 15 minutes. Call a physician.

**IF ON SKIN:** Wash thoroughly with soap and water. Remove and wash contaminated clothing before reuse.

**IF SWALLOWED:** Drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

**NOTE TO PHYSICIAN**

Irritable myocardial damage may contraindicate the use of gastric lavage. Measures against circulatory depression, respiratory depression, and convulsion may be needed.