

DEVERE

SANI KLEEN II

DISINFECTANT • CLEANER • FUNGICIDE
SANITIZER • MILDEWSTAY • DEODORANT
ACTIVE INGREDIENT

SANITIZATION (cont. from other panel)

thoroughly wetted surface for five minutes. Those surfaces that are heavily soiled must be pre-cleaned prior to sanitization (follow directions for cleaning).

To sanitize hard non-porous food contact surfaces such as equipment and utensils, mop or brush with a solution containing 1/4 ounce of SANI KLEEN II per gallon of water so as to maintain thoroughly wetted surface for 60 seconds. All surface must be thoroughly pre-cleaned with a good detergent followed by a potable water rinse (or follow directions for cleaning) prior to sanitization. Food contact surfaces must be rinsed with potable water before reuse. Do not reuse used solution.

STORAGE AND DISPOSAL

STORAGE: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. If plastic, do not reuse empty container.

DISPOSAL: Pesticide or rinsate that cannot be used or chemically reprocessed should be

ONLY FOR SALE TO, USE AND STORAGE
BY SERVICE PERSONS.

PRECAUTIONARY STATEMENTS

**HAZARDS TO HUMANS AND
DOMESTIC ANIMALS**

**DANGER: CAUSES EYE DAMAGE
CAUSES SEVERE SKIN IRRITATION, ESPECIALLY
TO BROKEN SKIN
HARMFUL IF SWALLOWED**

Do not get in eyes, on skin or on clothing.
Wear goggles or faceshield and rubber gloves when handling this
product.
Remove and wash contaminated clothing.
Do not take internally.

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Flush with plenty of water for at least 15 minutes.
Call a physician.

IF ON SKIN: Wash thoroughly with soap and water. Remove and
wash contaminated clothing before reuse.

IF SWALLOWED: Drink promptly a large quantity of milk, egg
whites, gelatin solution or if these are not available, drink large
quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric
lavage. Measures against circulatory shock, respiratory
depression and convulsion may be needed.

BEST DOCUMENT AVAILABLE

DIRECTIONS FOR USE

GENERAL CLASSIFICATION - It is a violation of federal law to use this product in a manner inconsistent with its labeling.

FOR CLEANING LIGHTLY SOILED AREAS - Surfaces to be cleaned should be mopped or brushed with a solution containing 1 ounce of SANI KLEEN II per gallon of water. Rinse cleaned area with fresh water.

FOR CLEANING HEAVILY SOILED AREAS - Mop or brush soiled surface with a solution containing 2 to 3 ounces of SANI KLEEN II per gallon of water. Rinse clean surface with fresh water.

GENERAL PURPOSE DISINFECTION - To disinfect walls, floors and other hard, non-porous surfaces in homes, schools, restaurants, restrooms and other public areas, mop or brush with a fresh solution containing 1 ounce of SANI KLEEN II per 1 gallon of water so as to maintain thoroughly wetted surface for 10 minutes. All soiled surfaces must be pre-cleaned prior to disinfection (follow directions for cleaning). Those surfaces which may later come in contact with food must be rinsed with clear water of drinking quality. Do not reuse used solution.

To disinfect hard, non-porous surfaces in locker rooms, shower stalls, walkways around pools and other areas against pathogenic fungi, mop or brush with a fresh solution containing 3/4 ounce of SANI KLEEN II per gallon of water so as to maintain thoroughly wetted surface for 10 minutes. This solution will destroy *Trichophyton interdigitale*, a causative agent of athlete's foot. All soiled surfaces must be pre-cleaned prior to disinfection (follow directions for cleaning). Do not reuse used solution.

To disinfect toilet bowls, flush toilet then force the bowl water over the trap. Add directly to bowl water 2 ounces of SANI KLEEN II. Swab or scrub all surfaces thoroughly including under the rim, and allow to remain wet for at least ten minutes before flushing toilet.

HOSPITAL DISINFECTION - To disinfect walls, floors, furniture and other hard, non-porous surfaces in hospitals and nursing homes, mop or brush with a fresh solution containing 2 ounces of SANI KLEEN II per gallon of water so as to maintain thoroughly wetted surface for 10 minutes. All soiled surfaces must be pre-cleaned prior to disinfection (follow directions for cleaning). Those surfaces that are heavily soiled with blood, fecal material or excreta should be thoroughly scrubbed with a solution containing 2 ounces of SANI KLEEN II per gallon of water, and then rinsed with a fresh solution containing 2 ounces of SANI KLEEN II per gallon of water before disinfection. Do not reuse used solution. This product has not been tested for effectiveness against *Mycobacterium tuberculosis* and must not be relied upon when tuberculocidal activity is intended by the user.

DEODORIZATION - Prior to deodorization of bathroom areas, garbage pails, diaper pails, food storage areas, kennel areas and other areas where obnoxious odors may develop, remove all garbage, refuse, food and/or animals. Pre-clean surfaces using pressurized water where possible before deodorization. To deodorize, mop or brush surfaces thoroughly with a solution containing 2 ounces of SANI KLEEN II per gallon of water.

MOLD AND MILDEW CONTROL - To control mold and mildew on previously cleaned glass, metal, plastic, tile and other hard non-porous surfaces, mop or brush with a fresh solution containing 2-1/2 ounces of SANI KLEEN II per gallon of water so as to thoroughly wet all surfaces. Allow to dry on surface. Repeat application weekly or when growth is visible. Do not reuse used solution.

SANITIZATION - To sanitize walls, floors and other hard non-porous, non-food contact surfaces in the presence of a moderate amount of organic soil, mop or brush with a solution containing 1/4 ounce of SANI KLEEN II per gallon of water so as to maintain

(cont. on other panel)