

2

Bio-Guard®

STA-CHLOR 60 CHLORINE CONCENTRATE


ACTIVE INGREDIENT:	66.52%
Trichloro-s-triazinetrione	
(available chlorine — 60%)	
INERT INGREDIENTS:	33.48%
TOTAL INGREDIENTS	100.00%

WARNING

KEEP OUT OF REACH OF CHILDREN
(SEE BACK PANEL
FOR ADDITIONAL PRECAUTIONS)

EPA Reg. No. 5185-85

Net Weight:



5185-85

When used as directed, this product is effective as a pool water disinfecting agent.

The filter system should be backwashed or properly cleaned and the pH adjusted between 7.2 and 7.6.

It is recommended that the pool be stabilized with 40-50 ppm trihydroxytriazine to nullify the dissipation effect of sunlight on chlorine. When using this product, both chlorine and trihydroxytriazine are being added to the water; however, it is necessary to initially stabilize pools with 40-50 ppm of trihydroxytriazine to effectively maintain the desired concentration of stabilizer.

DIRECTIONS

Use 4 ounces of Pool Chlorine Concentrate per 10,000 gallons of water as an initial treatment. Repeat daily until a residual of 1.0 to 3.0 ppm is established as determined by a chlorine test kit. The regular use of a test kit will aid in determining the amount of chlorine to be added for maintaining a residual of 1.0 to 3.0 ppm of available chlorine in the pool water. An average is 2-3 ounces of this chlorine concentrate per 10,000 gallons of water daily. The final demand for chlorine is to be established through use of a test kit and daily treatment adjusted accordingly.

The use of a quaternary ammonium type algicide will complement this total program of swimming pool maintenance.

If a need for superchlorination is indicated, use 1 pound of this concentrate per 10,000 gallons of water.



POISON



ANTIDOTE: Call a physician.

EXTERNAL: For skin and eyes, immediately flood with large amounts of water.

INTERNAL: Take large amounts of milk, followed by olive oil or cooking oil.

WARNING: Harmful or fatal if swallowed. May cause irritation to skin, nose and throat. Avoid contact with skin and eyes.

In case of contact, flush skin or eyes with cold water for at least 15 minutes. For eyes, get medical attention. Avoid breathing dust. **DO NOT TAKE INTERNALLY!**


Store in cool, dry, well ventilated areas. Keep containers closed when not in use. Contact with combustible matter may cause fire or decomposition. Keep away from heat or open flame. Oxidizing material. Will burn with the evolution of irritating or hazardous gases.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

KEEP OUT OF REACH OF CHILDREN

EPA Reg. No. 5185-85

Net Weight



5185-85

When used as directed, this product is effective as a pool water disinfecting agent.

The filter system should be backwashed or properly cleaned and the pH adjusted between 7.2 and 7.6.

It is recommended that the pool be stabilized with 40-50 ppms trihydroxytriazine to nullify the dissipation effect of sunlight on chlorine. When using this product, both chlorine and trihydroxytriazine are being added to the water; however, it is necessary to initially stabilize pools with 40-50 ppms of trihydroxytriazine to effectively maintain the desired concentration of stabilizer.



POISON

ANTIDOTE: Call a physician.

EXTERNAL: For skin and eyes, immediately flood with large amounts of water.

INTERNAL: Take large amounts of milk, followed by olive oil or cooking oil.

WARNING: Harmful . . . fatal if swallowed. May cause irritation to skin, nose and throat. Avoid contact with skin and eyes.

KEEP OUT OF REACH

EPA Reg. No. 5185-85