

Bio-Guard®

CHLORINE CONCENTRATE 750

ACTIVE INGREDIENTS:

Potassium dichloro-s-triazinetrione 76.14%

Trichloro-s-triazinetrione 16.63%

(Available chlorine—60%)

Total Active Ingredients 92.77%

INERT INGREDIENTS: 7.23%

WARNING

KEEP OUT OF REACH OF CHILDREN
(See back panel for additional precautions)

SOLD BY:

BIO-LAB, INC.
DECATUR, GEORGIA

ACCEPTED
SEP 1 1971
UNDER THE FEDERAL INSECTICIDE ACT
FUNGICIDE, HERBICIDE, AND
FOR ECOTOXICITY TESTS IN REGISTERED
ED UNITS 3185-95

U.S.D.A. Reg. No. 5185-95

Net Weight: _____

When used as directed, this product is effective as a pool water disinfecting agent.

The filter system should be backwashed or properly cleaned and the pH adjusted between 7.2 and 7.6.


It is recommended that the pool be stabilized with 50-60 ppm trihydroxytriazine to nullify the dissipation effect of sunlight on chlorine. When using this product, both chlorine and trihydroxytriazine are being added to the water; however, it is necessary to initially stabilize pools with 50-60 ppm of trihydroxytriazine to effectively maintain the desired concentration of stabilizer.

DIRECTIONS

Use 4 ounces of Pool Chlorine Concentrate per 10,000 gallons of water as an initial treatment. Repeat daily until a residual of 1.0 to 3.0 ppm of available chlorine is established. The regular use of a test kit will determine the amount of chlorine necessary for maintaining 1.0 to 3.0 ppm. An average is 2-3 ounces of this chlorine concentrate per 10,000 gallons of water daily. The final demand for chlorine is to be established through use of a test kit and daily treatment adjusted accordingly.

The use of a quaternary ammonium type algicide will complement this total program of swimming pool maintenance.

If a need for superchlorination is indicated, use 1 pound of this concentrate per 10,000 gallons of water.



Harmful if swallowed. May cause irritation of skin, eyes, nose and throat. Avoid contact with skin and eyes. Avoid breathing dust. **DO NOT TAKE INTERNALLY!** Store in well ventilated place. Keep container closed when not in use. Clean contaminated clothing before reuse.

ANTIDOTE: Call a physician.

External: For skin and eyes, immediately flood with large amounts of water.

Internal: Take large amounts of milk, followed by olive oil or cooking oil.

WARNING: Oxidizing material. Contact with heat, moisture or foreign materials may result in fire. Will burn with the evolution of irritating and hazardous gases.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

U.S.D.A. Reg. No. 5185-95

Net Weight: _____

PAPER SCREEN OF LABEL
SCREENED ON BOTTLE

Bio-Guard®

CHLORINE CONCENTRATE

750

ACTIVE INGREDIENTS:

Potassium dichloro-s-triazinetriene 76.14%
 Trichloro-s-triazinetriene 16.63%
 (Available chlorine—60%)

Total Active Ingredients 92.77%

INERT INGREDIENTS: 7.23%

WARNING

KEEP OUT OF REACH OF CHILDREN
 (See back panel for additional precautions)

SOLD BY:

BIO-LAB, INC.
 DECATUR, GEORGIA

ACCEPTED
 SEP 1 1971
 UNDER THE FEDERAL RESISTANCE
 FUNGICIDE AND HERBICIDE ACT
 FOR ECONOMIC POISON REGULATION
 FD UNL N. 2. 6105-95

U.S.D.A. Reg. No. 5185-95

Net Weight: _____

When used as directed, this product is effective as a pool water disinfecting agent.

The filter system should be backwashed or properly cleaned and the pH adjusted between 7.2 and 7.6.

It is recommended that the pool be stabilized with 50-60 ppm trihydroxytriazine to nullify the dissipation effect of sunlight on chlorine. When using this product, both chlorine and trihydroxytriazine are being added to the water; however, it is necessary to initially stabilize pools with 50-60 ppm of trihydroxytriazine to effectively maintain the desired concentration of stabilizer.

DIRECTIONS

Use 4 ounces of Pool Chlorine Concentrate per 10,000 gallons of water as an initial treatment. Repeat daily until a residual of 1.0 to 3.0 ppm of available chlorine is established. The regular use of a test kit will determine the amount of chlorine necessary for maintaining 1.0 to 3.0 ppm. An average is 2-3 ounces of this chlorine concentrate per 10,000 gallons of water daily. The final demand for chlorine is to be established through use of a test kit and daily treatment adjusted accordingly.

The use of a quaternary ammonium type algicide will complement this total program of swimming pool maintenance.

If a need for superchlorination is indicated, use 1 pound of this concentrate per 10,000 gallons of water.



Harmful if swallowed. May cause irritation of skin, eyes, nose and throat. Avoid contact with skin and eyes. Avoid breathing dust. **DO NOT TAKE INTERNALLY!** Store in well ventilated place. Keep container closed when not in use. Clean contaminated clothing before reuse.

ANTIDOTE: Call a physician.

External: For skin and eyes, immediately flood with large amounts of water.

Internal: Take large amounts of milk, followed by olive oil or cooking oil.

WARNING: Oxidizing material. Contact with heat, moisture or foreign materials may result in fire. Will burn with the evolution of irritating and hazardous gases.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

U.S.D.A. Reg. No. 5185-95

Net Weight: _____

PAPER SCREEN OF LABEL
 IS SCREENED ON BOTTLE