

5185-100 ACCEPTED May 21972

When used as directed this product is effective as a pool water disinfecting agent.

The filter system should be backwashed or properly cleaned and the pH adjusted between 7.4 and 7.6.

It is recommended that the pool be stabilized with 40-50 ppm trihydroxy triazine to nullify the dissipation effect of sunlight on chlorine. When using this product, both chlorine and trihydroxy triazine are being added to the water; however, it is necessary to initially stabilize pools with 40-50 ppm of trihydroxy triazine to effectively maintain the desired concentration of stabilizer.

DIRECTIONS

Use 3 ounces of Pool Chlorine Concentrate per 10,000 gallons of water as an initial treatment. Repeat until a residual of 1.0 to 2.0 ppm of available chlorine is established as determined by test kit. A test kit must be used regularly to determine when it is necessary to add another dose of chlorine concentrate to maintain the residual of 1.0 to 2.0 ppm of available chlorine in the pool water. An average is 2-3 ounces of this chlorine concentrate per 10,000 gallons of water daily.



DANGER!

OXIDIZING MATERIAL. CONTACT WITH HEAT, MOISTURE OR FOREIGN MATERIALS MAY RESULT IN FIRE. WILL BURN WITH THE EVOLUTION OF IRRITATING AND HAZARDOUS GASES.

Store in a cool dry place. Keep away from heat or open flame. Avoid contact with skin and eyes. Avoid breathing dust. DO NOT TAKE INTERNALLY! May be fatal if swallowed.

This product is toxic to fish. Treated effluent should not be discharged where it will drain into lakes, streams, ponds, or public water. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

ANTIDOTE: Call a physician.

EXTERNAL: For skin and eyes, immediately flood with large amounts of water.

INTERNAL: Take large amounts of milk, followed by olive oil or cooking oil.

E.P.A. Reg. No.

KEEP OUT OF THE REACH OF CHILDREN

NET WEIGHT 10 POUNDS

35