

# Pool-Guard®

**ACCEPTED**  
June 29, 1977  
UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTERED  
ED UNDER NO. 5135-105

## 60-65 Chlorine CONCENTRATE

Active Ingredients:	
Trichloro-s-triazinetrione	72.10%
(Available chlorine--65%)	
Inert Ingredients:	27.90%
Total Ingredients	100.00%

**WARNING**  
**KEEP OUT OF REACH OF CHILDREN**  
See side panel for additional precautions.

U.S.D.A. Reg. No. 5135-105      Net Weight:

Sold by  
**BIO-LAB, INC.**  
DECATUR, GEORGIA

NEAREST SUPERMART REPRESENTS APPROXIMATELY 1 OZ.

When used directed, this product is effective as a pool water disinfecting agent.

The filter system should be backwashed or properly cleaned and the pH adjusted between 7.2 and 7.6.

It is recommended that the pool be stabilized with 40-50 ppm trihydroxy triazine to nullify the dissipation effect of sunlight on chlorine. When using this product, both chlorine and trihydroxy triazine are being added to the water; however, it is necessary to initially stabilize pools with 40-50 ppm of trihydroxy triazine to effectively maintain the desired concentration of stabilizer.

### DIRECTIONS

Use 4 ounces of Pool Chlorine Concentrate per 10,000 gallons of water as an initial treatment. Repeat daily until a residual of 1.0 to 3.0 ppm of available chlorine is established. The regular use of a test kit will aid in determining the amount of chlorine necessary for maintaining 1.0 to 3.0 ppm. An average is 2-3 ounces of this chlorine concentrate per 10,000 gallons of water daily. The final demand for chlorine is to be established through use of a test kit and daily treatment adjusted accordingly.

The use of a quaternary ammonium type algicide will complement this total program of swimming pool maintenance. If a need for superchlorination is indicated, use 1 pound of this concentrate per 10,000 gallons of water.

### POISON

Harmful if swallowed. **DO NOT TAKE INTERNALLY!** Avoid contact with skin and eyes. Avoid breathing dust. Keep cover closed when not in use. Protect from moisture contamination.

**Antidote:** Call a physician.  
External: For skin and eyes, immediately flood with large amounts of water. For eyes get immediate medical attention.  
Internal: Take large amounts of milk, followed by olive oil or cooking oil.

### KEEP OUT OF REACH OF CHILDREN

**WARNING:** Oxidizing material. Contact with heat, moisture or foreign materials may result in fire. Will burn with the evolution of irritating and toxic gases. Store in cool dry place. Keep away from heat or open flame.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

DO NOT REUSE  
OR BURN

# Bio-Guard®

ACCEPTED  
June 29, 1971  
UNDER THE FEDERAL INSECTICIDE, FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON RESISTANCE NO. 5185-105

## 60-65 Chlorine CONCENTRATE

EACH GRADUATION REPRESENTS APPROXIMATELY 1 OZ.

Active Ingredients:		
Trichloro-s-triazinirone (Available chlorine—65%)	72.10%	
Inert Ingredients:	27.90%	
Total Ingredients	100.00%	

**WARNING**  
KEEP OUT OF REACH OF CHILDREN  
See side panel for additional precautions.

U.S.D.A. Reg. No. 5185-105 Net Weight

Sold by  
**BIO-LAB, INC.**  
DECATUR, GEORGIA

BEST DOCUMENT AVAILABLE

WITH SCREEN OF LAST WHICH IS SCREENED ON

When used as directed, this product is effective as a pool water disinfecting agent.

The filter system should be backwashed or properly cleaned and the pH adjusted between 7.2 and 7.6.

It is recommended that the pool be stabilized with 40-50 ppm trihydroxy triazine to nullify the dissipation effect of sunlight on chlorine. When using this product, both chlorine and trihydroxy triazine are being added to the water; however, it is necessary to initially stabilize pools with 40-50 ppm of trihydroxy triazine to effectively maintain the desired concentration of stabilizer.

### DIRECTIONS

Use 4 ounces of Pool Chlorine Concentrate per 10,000 gallons of water as an initial treatment. Repeat daily until a residual of 1.0 to 3.0 ppm of available chlorine is established. The regular use of a test kit will aid in determining the amount of chlorine necessary for maintaining 1.0 to 3.0 ppm. An average is 2-3 ounces of this chlorine concentrate per 10,000 gallons of water daily. The final demand for chlorine is to be established through use of a test kit and daily treatment adjusted accordingly.

The use of a quaternary ammonium type algicide will complement this total program of swimming pool maintenance. If a need for superchlorination is indicated, use 1 pound of this concentrate per 10,000 gallons of water.

### POISON

Harmful if swallowed. DO NOT TAKE INTERNALLY! Avoid contact with skin and eyes. Avoid breathing dust. Keep cover closed when not in use. Protect from moisture contamination.

Antidote: Call a physician.  
External: For skin and eyes, immediately flood with large amounts of water. For eyes get immediate medical attention.  
Internal: Take large amounts of milk, followed by olive oil or cooking oil.

### KEEP OUT OF REACH OF CHILDREN

WARNING: Oxidizing material. Contact with heat, moisture or foreign materials may result in fire. Will burn with the evolution of irritating and hazardous gases. Store in cool dry place. Keep away from heat or open flame.

Do not reuse empty container. Destroy it by burning with burning. Stay away from smoke or fumes.

