

3

APR 21 1971
5185-135

kleer pool

SWIMMING POOL WATER
TREATMENT PROGRAM

CHLORINATED STABILIZER

ACTIVE INGREDIENTS:	
Sodium dichloro-s-triazinetrione	38.30%
INERT INGREDIENTS:	61.70%
TOTAL:	100.00%

WARNING

KEEP OUT OF REACH OF CHILDREN
See additional warnings on back panel

BIO-LAB, INC. / DECATUR, GA.

5185-135
ACCEPTED

Aug. 9, 1971

U.S. DEPARTMENT OF AGRICULTURE
FOOD AND NUTRITION ASSISTANCE
PROGRAM
FEDERAL REGISTER
U.S. GOVERNMENT PRINTING OFFICE

ACTIVE INGREDIENTS:

Sodium dichloro-s-triazinetrione 38.30%

INERT INGREDIENTS:* 61.70%

TOTAL: 100.00%

*Includes Trihydroxy Triazine (chlorine stabilizer).

This product is formulated as an initial chemical treatment for swimming pool water. When used as directed, it supplies adequate available chlorine to satisfy the initial demand in pool water from plant purification sources, and also supplies 50 parts per million of Trihydroxy triazine (chlorine stabilizer). It also contains a buffering agent to aid in the control of pH of pool water, and a sequestering agent to aid in the prevention of precipitation of salts of metallic ions.

DIRECTIONS

Backwash the filter system before adding this product to the pool. Establish a pool water pH of between 7.4 and 7.6. Then add one bag (3½ lbs.) of this product per 5000 gallons of water. Add chemical at night, applying through the skimmer. Maintain pool water circulation continuously.

When the chemical reaches the filter system there may be a temporary increase in filter pressure, which will dissipate as the chemical is dissolved. Do not backwash the filter for 24 hours after application of this chemical.

There will be an initial high chlorine reading and the pool should not be used until residual chlorine is at a normal level of 1.0-3.0 parts per million as determined by test kit. Subsequent treatment with chlorine should be initiated to maintain a chlorine residual of 1.0-3.0 parts per million as determined by test kit.

U.S.D.A. Reg. No. 5185-135

Net Wt. 3.5 lbs.

ANTIDOTE: Call a physician.

EXTERNAL: For skin and eyes, immediately flood with large amounts of water.

INTERNAL: Take large amounts of milk, followed by olive oil or cooking oil.

KEEP OUT OF THE REACH OF CHILDREN

WARNING: OXIDIZING MATERIAL. CONTACT WITH HEAT, MOISTURE OR FOREIGN MATERIALS MAY RESULT IN FIRE, WILL BURN WITH THE EVOLUTION OF IRRITATING AND HAZARDOUS GASES.

Store in a cool dry place. Keep away from heat or open flame. Avoid contact with skin and eyes. Avoid breathing dust. **DO NOT TAKE INTERNALLY! DO NOT REUSE CONTAINER. DESTROY WHEN EMPTY.**

**KLE
POC**
SWIMMING
TREATMENT

CHLORINATED S

ACTIVE INGREDIENTS:
Sodium dichloro-s-tri
INERT INGREDIENTS:*

WARN
KEEP OUT OF REAC
See additional warnir

BIO-LAB, INC. / E

ACTIVE INGREDIENTS:**Sodium dichloro-s-triazinetriene 38.30%****INERT INGREDIENTS:* 61.70%****TOTAL: 100.00%**

*Includes Trihydroxy Triazine (chlorine stabilizer).

This product is formulated as an initial chemical treatment for swimming pool water. When used as directed, it supplies adequate available chlorine to satisfy the initial demand in pool water from plant purification sources, and also supplies 50 parts per million of Trihydroxy triazine (chlorine stabilizer). It also contains a buffering agent to aid in the control of pH of pool water, and a sequestering agent to aid in the prevention of precipitation of salts of metallic ions.

DIRECTIONS

Backwash the filter system before adding this product to the pool. Establish a pool water pH of between 7.4 and 7.6. Then add one bag (3½ lbs.) of this product per 5000 gallons of water. Add chemical at night, applying through the skimmer. Maintain pool water circulation continuously.

When the chemical reaches the filter system there may be a temporary increase in filter pressure, which will dissipate as the chemical is dissolved. Do not backwash the filter for 24 hours after application of this chemical.

There will be an initial high chlorine reading and the pool should not be used until residual chlorine is at a normal level of 1.0-3.0 parts per million as determined by test kit. Subsequent treatment with chlorine should be initiated to maintain a chlorine residual of 1.0-3.0 parts per million as determined by test kit.

U.S.D.A. Reg. No. 5185-135**Net Wt. 3.5 lbs.****ANTIDOTE:** Call a physician.**EXTERNAL:** For skin and eyes, immediately flood with large amounts of water.**INTERNAL:** Take large amounts of milk, followed by olive oil or cooking oil.**KEEP OUT OF THE REACH OF CHILDREN****WARNING:** OXIDIZING MATERIAL. CONTACT WITH HEAT, MOISTURE OR FOREIGN MATERIALS MAY RESULT IN FIRE, WILL BURN WITH THE EVOLUTION OF IRRITATING AND HAZARDOUS GASES.

Store in a cool dry place. Keep away from heat or open flame. Avoid contact with skin and eyes. Avoid breathing dust. **DO NOT TAKE INTERNALLY!**
DO NOT REUSE CONTAINER. DESTROY WHEN EMPTY.