

This product is effective as a pool water disinfecting agent when used as directed.

 q_{i}

. TICI (C

JULIA REGISTERA

a Constant

It is recommended that the pool be stabilized with 40-50 ppm trihydroxy triazine to nullify the dissipation effect of sunlight on chlorine. When using this product, both chlorine and trihydroxy triazine are being added to the water; however, it is necessary to initially stabilize puols with 40-50 ppm of trihydroxy triazine to effectively maintain the desired concentration of stabilizer.

DIRECTIONS

The filter system should be backwashed or properly cleaned and the pH adjusted between 7.4 and 7.6. Before using this product, it is recommended that the pool be super chlorinated so that a chlorine residual of between 1.0 and 2.0 ppm is obtained.

These tablets may be used in various types of proportioning mechanisms. Adjust whatever feeding mechanism that might be used to assure a constant treatment level of 1.0-2.0 ppm available chloring in the pool water (normally 1-2 ounces of this product per 10,000 gallons daily). (Four tablets is approximately one ounce.)

The regular use of a test kit will determine if further adjustments may be necessary to maintain the recommended level of available chlorine in the pool water.

CAUTION: DO NOT MIX OTHER CHEMICALS WITH THIS PRODUCT, AS A DANGEROUS REACTION MAY OCCUR.

DANGER

KEEP OUT OF THE REACH OF CHILDREN

Oxidizing material. Contact with heat, moisture or foreign materials may result in fire. Will burn with the evolution of irritating and hazardous gases.

Store in a cool dry place. Keep away from heat or open flame. Avoid contact with skin and eyes. Avoid breathing dust. DO NOT TAKE INTERNALLY! DO NOT REUSE CONTAINER. Rinse thoroughly with water and discard.

Antidote: Call a physician.

External: For skin and eyes, immediately flood with large amounts of water.

Internal Take large amounts of milk, followed by alive all or cooking ail.

NET CONTENTS: MANUFACTURED BY: BIO-LAS, INC., DECATUR, GEORGIA

