

FIRST AID


In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, get medical attention.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Call a physician immediately.

FOR INDUSTRIAL USE ONLY

EPA Est. No. 4140
CA-1 IN-1 NJ-1 PR-1 TX-1

TO REORDER
Call TOLL-FREE: 1-800-527-9921
In Texas: 1-800-442-7950



LEFT PANEL

KEM MANUFACTURING

GERMEX AEROSOL

HOSPITAL DISINFECTANT DEODORANT

Not Recommended for use in conjunction with
other disinfectants or deodorants.
Read label for complete directions of use.

ACTIVE INGREDIENTS:

Ortho-phenylphenol.....	0.10%
Para-tertiary-amylphenol.....	0.05%
Inert Ingredients.....	<u>99.85%</u>
	100.00%

EPA REG. NO. 5197-50

EPA Est. 4140-TX-1

WARNING

KEEP OUT OF REACH OF CHILDREN

FLAMMABLE MIXTURE

CONTENTS UNDER PRESSURE

See side panel for additional precautions.

NET CONTENTS: 13OZ./369G.

KEM MANUFACTURING

A Division of Systems General, Inc.

P.O. Box 152170

Irving, TX 75015-2170


CENTER PANEL

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

This product will control the following organisms: Aspergillus Niger, Escherichia Coli, Pseudomonas Aeruginosa, Salmonella Choleraesuis, Salmonella Typhosa, Staphylococcus Aureus, Trichophyton Interdigitale, Tubercle Bacilli on environmental surfaces.

For Disinfecting: Hold dispenser upright about 6 to 8 inches from pre-cleaned surface to be disinfected and spray 3 seconds until surface is wet.



For Deodorizing: Hold dispenser upright-point the opening away from face. Spray into air towards the center of the room for about 4 seconds.

For Mildew Control on Hard Surfaces: Clean surfaces before application. Spray surface until wet. Repeat application every week to maintain control.


This powerful hospital disinfectant and deodorant acts quickly on many germ-laden surfaces, such as: refuse cans, toilet seats, urinals, laundry chutes, wheel chairs, air ducts and light switches after the surface has been properly cleaned. Reduces the danger of transmission of many infectious organisms from environmental surfaces. Effective on odors from many sources in hospitals, nursing homes, ambulances.

NOTE: Do not use on rayon fabrics.

RIGHT PANEL

WARNING

Effects of Overexposure: **Eyes:** Contact may cause stinging or burning. **Skin:** Contact may cause drying of skin, cracking with irritation. **Inhalation:** May cause irritation and burning sensation of lips, mouth and throat, gastro-intestinal irritation, stomach cramps, nausea and vomiting.



LEFT PANEL

KEM MANUFACTURING

GERMEX AEROSOL

HOSPITAL DISINFECTANT DEODORANT

ACTIVE INGREDIENTS:

Ortho-phenylphenol.....	0.10%
Para-tertiary-amylphenol.....	0.05%
Inert Ingredients.....	99.85%
	<u>100.00%</u>

EPA REG. NO. 5197-50

EPA Est. 4140-TX-1

WARNING

KEEP OUT OF REACH OF CHILDREN

FLAMMABLE MIXTURE

CONTENTS UNDER PRESSURE

See side panel for additional precautions.

NET CONTENTS: 13OZ./369G.

KEM MANUFACTURING

A Division of Systems General. Inc.

P.O. Box 152170

Irving, TX 75015-2170

CENTER PANEL

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

This product will control the following organisms: Aspergillus Niger, Escherichia Coli, Pseudomonas Aeruginosa, Salmonella Choleraesuis, Salmonella Typhosa, Staphylococcus Aureus, Trichophyton Interdigitale, Tubercle Bacilli on environmental surfaces.

For Disinfecting: Hold dispenser upright about 6 to 8 inches from pre-cleaned surface to be disinfected and spray 3 seconds until surface is wet.

Not Reviewed. Registrant claims to be in accordance with EPA's Pesticide Product Guidance for Phase 2 Disinfectants, Page 2.12, Active to Inert Change in Status.

For Deodorizing: Hold dispenser upright-point the opening away from face. Spray into air towards the center of the room for about 4 seconds.

For Mildew Control on Hard Surfaces: Clean surfaces before application. Spray surface until wet. Repeat application every week to maintain control.


This powerful hospital disinfectant and deodorant acts quickly on many germ-laden surfaces, such as: refuse cans, toilet seats, urinals, laundry chutes, wheel chairs, air ducts and light switches after the surface has been properly cleaned. Reduces the danger of transmission of many infectious organisms from environmental surfaces. Effective on odors from many sources in hospitals, nursing homes, ambulances.

NOTE: Do not use on rayon fabrics.

RIGHT PANEL

WARNING

Effects of Overexposure: **Eyes:** Contact may cause stinging or burning. **Skin:** Contact may cause drying of skin, cracking with irritation. **Inhalation:** May cause irritation and burning sensation of lips, mouth and throat, gastro-intestinal irritation, stomach cramps, nausea and vomiting.



Emergency First Aid Procedures: **Eyes:** Immediately flush eyes with running water for at least 15 minutes. If pain and/or irritation persists, get medical attention. **Skin:** Remove contaminated clothing immediately and flush affected area with water for at least 15 minutes. If pain or irritation persists, get medical attention. Launder clothing before reusing. **Inhalation:** Move victim to fresh air. **Ingestion:** If swallowed, induce vomiting by giving a glass of salt water, touching fingers to back of throat, or by giving 30 cc (2 Tbs) Syrup of Ipecac. Keep victim warm and quiet. Get immediate medical attention. Do not give anything by mouth to an unconscious person.

Avoid contact with food. Rinse with water anything that will come in contact with baby's mouth. **KEEP OUT OF REACH OF CHILDREN.**

Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130F may cause busting.

NOT FOR RESALE
FOR INDUSTRIAL USE ONLY

TO REORDER
Call TOLL-FREE: 1-800-527-9921
In Texas: 1-800-442-7950

