

FOAMEX

For Post Harvest washing of Citrus Fruit
and control of Blue and Green Mold.

DIRECTIONS: FoameX is supplied as a concentrate and must be diluted prior to use. Mix 1 part FoameX concentrate with 9 parts water and apply as a foam over the fruit as it travels through a brush washer. Apply sufficient material to thoroughly cover the fruit. An exposure time of 20 seconds is recommended. Brushing in excess of 1 minute may result in burning of the fruit peel. Treatment must be followed by thorough fresh water rinse.

ACTIVE INGREDIENTS:

Sodium Ortho Phenylphenate, anhydrous.....14.52%

INERT INGREDIENTS:85.48%

CAUTION! KEEP OUT OF REACH OF CHILDREN.

Treated effluent waters may be harmful to fish and other aquatic life.

HARMFUL IF SWALLOWED

CORROSIVE LIQUID —

IF SPILLED ON SKIN OR CLOTHING, FLUSH WITH FRESH WATER.

When empty, replace bungs in barrel and return to Pomona.

E.P.A. Reg. No. 5202-12

Net Contents: 55 Gallons

BROGDEX
COMPANY®

Pomona, California

Form 101-68, Aug. 73

ACCEPTED

May 3, 1974
FEDERAL INSECTICIDE
AND RODENTICIDE ACT
NOMINATED FOR REGISTRATION
EPA REG. NO. 5202-12
EPA REG. NO. 5202-12