

**Precautionary  
Statements**

Hazards to Humans and  
Domestic Animals

**CAUTION**

HARMFUL IF  
SWALLOWED

Avoid contact with skin, eyes or  
clothing

**FIRST AID**

Avoid contact with skin, eyes or clothing. In  
case of contact immediately flush eyes or skin  
with plenty of water. Get medical attention if  
irritation persists.

**ENVIRONMENTAL  
HAZARDS**

The product is toxic to fish. Do not apply  
directly to water. Do not contaminate water  
by cleaning of equipment or disposal of  
waters. Do not dump where surface or sub-  
surface waters may become polluted.

**DIRECTIONS  
FOR USE**

In accordance with Federal law, do not use this product in a manner inconsistent with its labeling.

**STORAGE AND  
DISPOSAL**

Do not contaminate water,  
food or feed by storage or  
disposal.

**STORAGE:** Keep product from freezing.  
Satisfactory utility is expected  
after thawing and mixing of product.  
Openumping is prohibited.

**DISPOSAL:** Pesticide, spray mix-  
ture or residue that cannot be used  
according to label instructions must  
be disposed of according to applicable  
Federal, State or Local procedures.

**CONTAINERS:** Triple rinse for equiv-  
alent and offer for recycling or  
reconditioning. Do not dispose of in  
a sanitary landfill or by other  
approved state and local procedures.  
Plastic containers may be incinerated  
if permitted by state and local  
authorities.

BY THE MANUFACTURER  
FOR THE MANUFACTURER

# BROGDEX®

## 593-F

A postharvest fungicidal treatment for  
citrus fruits, apples and pears.

**CROP DIRECTIONS**

Brogdex 593-F does not have to  
be agitated if used before the  
date stamped below:

Use Before \_\_\_\_\_ or Agitate

If not used before the above date  
the container must be thoroughly  
mixed. Mixing should be repeated  
every 30 days or gentle agitation  
on a continuous basis may be used.

**CITRUS FRUITS:** (Postharvest) Blue  
Mold, Green Mold and Stem-end rot  
Dilute one gallon of Brogdex 593-F  
to 30 gallons with water for a  
1000 ppm treating suspension.  
Treat clean fruit by dip, spray  
or flood application. Drain treated  
fruit thoroughly, do not rinse  
treated fruit.

**APPLES:** (Postharvest) Blue Mold,  
Gray Mold, Bull's eye rot -  
Dilute one gallon Brogdex 593-F  
to 30 gallons with water for a  
1000 ppm treating suspension.  
Treat sound fruit by dip, spray  
or flood application. Treat  
fruit before and after storage  
for maximum disease control. Do  
not treat for more than three  
minutes per application. Drain  
fruit thoroughly and do not  
rinse after treatment.

**PEARS:** (Postharvest) Blue Mold,  
Gray Mold, Bull's eye rot -  
Dilute one gallon Brogdex 593-F  
to 30 gallons with water for a  
1000 ppm treating suspension.  
Treat sound fruit only once, either  
before or after storage, for not  
more than three minutes. Treat by  
dip, spray or flood. Drain fruit  
thoroughly and do not rinse  
after treatment.

**NOTE:** Fungi may develop tolerance  
to Brogdex 593-F. If such tolerance  
(or resistance) is suspected,  
switching to a non-benzimidazole  
fungicide is advised. Check with  
your supplier if confirmatory  
tests are desired.

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

See Side Panel For Additional Precautionary Statements

# BROGDEX® COMPANY

1441 West Second Street, Pomona, California

Net Contents \_\_\_\_\_ U.S. Gallons

EPA Reg. No. 8202-

EPA Est. 8202-CA-1

ACCEPTED

DRP 3 1982