

3200-11
ACCEPTED
 REV 4/1/70
 UNDER THE
 FUNGICIDE
 FOR THE
 ED USE



CAUTION
 KEEP OUT
 OF REACH OF
 CHILDREN

SEE SIDE
 PANEL FOR
 ADDITIONAL
 CAUTIONS

**JAPANESE BEETLE GRUBS
 SOD WEBWORMS • ANTS
 TERMITES**

ACTIVE INGREDIENT:
 Technical Chlordane 50.00%
INERT INGREDIENTS: 50.00%
 *Equivalent to 30% Octachloro-4, 7--Methanotetrahy-
 droindane and 20% related compounds on a weight
 basis

CAUTION

KEEP OUT OF REACH OF CHILDREN and PETS.
 CHLORDANE is toxic to humans and warm blood-
 ed animals.

- Avoid ingestion. Harmful if swallowed.
- Wash hands and face thoroughly with soap and water after use, or before eating and smoking.
- Avoid contact.
- Avoid contamination of foodstuffs and feed.
- Do not apply to food crops after edible parts begin to form.
- Do not use on dairy animals.
- Do not use on forage crops to be fed dairy animals or animals being finished for slaughter.
- Do not use in dairy barns or poultry houses.
- Rinse empty container thoroughly with water and dispose of it.

FISH and WILDLIFE CAUTION

Toxic to fish and wildlife. Birds feeding on treated areas may be killed. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not contaminate any body of water nor apply when conditions favor runoff from area treated.

ANTIDOTE

If swallowed, give one teaspoon of baking soda dissolved in a glass of warm water and repeat every 15 minutes. CALL A PHYSICIAN.

AGRICULTURAL PRODUCTS CO.
 WEST KINGSTON, R. I.

NET CONTENTS

LAWNS

To control Japanese Beetles, Asiatic Beetles, Ground Squirrels, apply APCO CHLORDANE 50% WETTABLE at 2 tablespoons per 1000 square feet. For Ants, Earthworms, Sod Webworms, apply 1/2 gallon of water per 500 square feet. For Japanese Beetles, Sod Webworms, apply 50% Wettable at 2 tablespoons per 1000 square feet. Wash the lawn with water after application. Wash hands with soap and water after use.

FOR USE ON SUBTERRANEAN

SOIL TERMITES
 Effective against termites, ants, and other insects. Apply to soil in the form of a bait.

FRENCH CRICKETS
 To be effective, apply to soil in the form of a bait. Apply to soil in the form of a bait. Apply to soil in the form of a bait.

IT IS IMPORTANT TO APPLY UNIFORMELY TO THE DEPTH OF CHEMICAL ACTION.
 After application, water the soil to help the chemical penetrate.

CAUTION

KEEP OUT OF REACH OF CHILDREN and PETS.
Chlordane is toxic to fish and aquatic life.

Do not use if swallowed.
Wash hands thoroughly with soap and water after use. Do not eat, drink, or smoke while using.

Do not use on food crops.
Do not use on food, feed, or bedding.
Do not use on children's toys or clothing.

Do not use on dairy animals.
Do not use on dairy animals to be fed dairy products.
Do not use on animals for slaughter.

Do not use in dairy barns or poultry houses.
Rinse empty containers thoroughly with water and dispose of it.

FISH AND WILDLIFE CAUTION

Toxic to fish and wildlife. Birds feeding on treated areas may be killed. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not contaminate any body of water nor apply within 100 feet of surface runoff from area treated.

ANTIDOTE

If swallowed, give oral solution of baking soda or saturated solution of sodium bicarbonate and repeat until vomit is clear. CALL A PHYSICIAN.

AGRICULTURAL PRODUCTS CO
WEST KINGSTON, R. I.



CAUTION
KEEP OUT
OF REACH OF
CHILDREN

SEE SIDE
PANEL FOR
ADDITIONAL
CAUTIONS



ACTIVE INGREDIENT:
Technical Chlordane 50.00%
INERT INGREDIENTS: 50.00%
*Equivalent to 30% Octachloro-1,7-Methano-1,4-dioxin and 20% related compounds in a weight basis.

NET CONTENTS ONE POUND
SUGGESTED USE

LAWNS

To control Japanese Beetle, European Chalcid Beetle Grubs, Mole Crickets, and W. P. apply APCO CHLORDANE 50% W. P. in water per 100 sq. ft. of lawn or turf.

For Ants, Earwigs, and Sod Webworms, apply 1/2 lb. per 100 sq. ft. of lawn.

For control of Japanese Beetle, European Chalcid Beetle Grubs, Mole Crickets, and W. P. apply 1/2 lb. per 100 sq. ft. of lawn. To wash the insecticide off the lawn, use W. P. per 100 sq. ft. of lawn.

FOR CONTROL OF SUBTERRANEAN TERMITES

SOIL APPLICATION:
CHLORDANE 50% W. P. is applied to the soil in a 1/2 inch layer, 1/2 inch deep, in the fourth row of the trench, and other rows as needed.

TRENCHING PROCEDURE:
To dig a trench, use a trenching machine or hand-dug trench. The trench should be 1/2 inch deep and 1/2 inch wide. The soil in the trench should be moist. APCO CHLORDANE 50% W. P. is applied to the soil in a 1/2 inch layer, 1/2 inch deep, in the fourth row of the trench, and other rows as needed.

IT IS IMPORTANT TO HAVE THE CHEMICAL UNIFORMLY DISTRIBUTED TO THE DEPTH OF THE SUBTERRANEAN CHEMICAL BARRIER.

After the trenching is completed, the soil should be replaced and the trench filled with soil.

