

DIRECTIONS:

Apply to leaves. Do not wet plants. Do not use on plants with frost damage. Do not use on plants with insecticide resistance.

Lawns

Apply to lawn. Do not use on plants with frost damage. Do not use on plants with insecticide resistance.

Do not use on plants with frost damage. Do not use on plants with insecticide resistance.

Cabbage Looper, Harlequin Cabbage, Imported Cabbage Worm, Cabbage Root Fly, etc. per 100 square feet.

Cabbage

Cabbage Moth, etc. per 100 square feet. Apply to plants after they are established.

Strawberry


Do not apply to strawberries after the blossoms open.

Do not apply to strawberries after the blossoms open.

AGRICULTURAL PRODUCTS CO.
WEST HAVEN, CONNECTICUT



ACCEPTED
1-30-71
5253-13



CAUTION
KEEP OUT OF REACH
OF CHILDREN

SEE SIDE
PANEL FOR
ADDITIONAL
CAUTIONS

**JAPANESE BEETLE GRUBS.
SOD. WEBWORMS • ANTS**

ACTIVE INGREDIENTS
CHLORDANE 6%
INERT INGREDIENTS 94%

NET WEIGHT 10 LBS. ROUND

NET WEIGHT 10 LBS. ROUND

Do not use on plants with frost damage. Do not use on plants with insecticide resistance.

CAUTION

KEEP OUT OF REACH OF CHILDREN AND DOMESTIC ANIMALS.

Do not use on plants with frost damage. Do not use on plants with insecticide resistance.

Do not use on plants with frost damage. Do not use on plants with insecticide resistance.